# <sup>by</sup> Impact of a multifaceted and multidisciplinary intervention on pain, agitation and delirium management in an intensive care unit: an experience of a Canadian community hospital in conducting a quality improvement project

Zechen Ma <sup>1</sup>, <sup>1</sup> Mercedes Camargo Penuela,<sup>2,3</sup> Madelyn Law,<sup>3</sup> Divya Joshi,<sup>4</sup> Han-Oh Chung,<sup>2,5</sup> Joyce Nga Hei Lam,<sup>2</sup> Jennifer LY Tsang <sup>1</sup>,<sup>5,6</sup>

## ABSTRACT

**To cite:** Ma Z, Camargo Penuela M, Law M, *et al.* Impact of a multifaceted and multidisciplinary intervention on pain, agitation and delirium management in an intensive care unit: an experience of a Canadian community hospital in conducting a quality improvement project. *BMJ Open Quality* 2021;**10**:e001305. doi:10.1136/ bmjoq-2020-001305

Additional supplemental material is published online only. To view, please visit the journal online (http://dx.doi.org/10. 1136/bmjoq-2020-001305).

Received 15 December 2020 Accepted 23 November 2021

#### Check for updates

© Author(s) (or their employer(s)) 2021. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

For numbered affiliations see end of article.

#### **Correspondence to**

Dr Jennifer LY Tsang; jennifer.tsang@mail.utoronto.ca

Background Clinical guidelines suggest that routine assessment, treatment, and prevention of pain, agitation, and delirium (PAD) is essential to improving patient outcomes as delirium is associated with increased mortality and morbidity. Despite the well-established improvements on patient outcomes, adherence to PAD guidelines is poor in community intensive care units (ICU). This guality improvement (QI) project aims to evaluate the impact of a multifaceted and multidisciplinary intervention on PAD management in a Canadian community ICU and to describe the experience of a Canadian community hospital in conducting a QI project. Methods A ten-member PAD advisory committee was formed to develop and implement the intervention. The intervention consisted of a multidisciplinary rounds script, poster, interviews, visual reminders, educational modules, pamphlet and video. The 4-week intervention targeted nurses, family members, physicians, and the multidisciplinary team. An uncontrolled, before-andafter study methodology was used. Adherence to PAD assessment guidelines by nurses was measured over a 6-week pre-intervention and over a 6-week postintervention periods.

**Results** Data on 430 and 406 patient-days (PD) were available for analysis during the pre- and post-intervention periods, respectively. The intervention did not improve the proportion of PD with guideline compliance to the assessment of pain (23.4% vs. 22.4%, p=0.80), agitation (42.9% vs. 38.9%, p=0.28), nor delirium (35.2% vs. 29.6%, p=0.10) by nurses.

**Discussion** The implementation of a multifaceted and multidisciplinary intervention on PAD assessment did not result in significant improvements in guideline adherence in a community ICU. Barriers to knowledge translation are apparent at multiple levels including the personal level (low completion rates on educational modules), interventional level (under-collection of data), and organisational level (coinciding with hospital accreditation education). Our next steps include reintroduction of education modules using organisation approved platforms, updating existing ICU policy, updating admission order sets, and conducting audit and feedback.

## **INTRODUCTION**

Most patients admitted to the intensive care unit (ICU) experience pain even at rest, which can stem from surgery, trauma, burns, cancer and procedures.<sup>T</sup> The incidence of pain is high in both surgical-trauma patients (52%) as well as medical patients (50%) in the ICU.<sup>2</sup> Not only does pain affect patients for their duration in the ICU, the memory of unrelieved pain persists post-ICU discharge for many years and can be distressing. In addition, at least 71% of patients develop agitation in the ICU determined by bedside clinical judgement.3 Disease-induced and iatrogenic delirium can affect 32%-80% of critically ill patients in the ICU.<sup>4</sup> This in turn is an important independent negative predictor for several outcomes, including a threefold increase in 6-month mortality, extra days on mechanical ventilation, increased hospital length of stay,<sup>5</sup> and long-term cognitive impairment consistent with a dementialike state.<sup>467</sup>

In response to addressing pain, agitation and delirium (PAD) experienced by patients in the ICU, the Society of Critical Care Medicine published clinical practice guidelines for the assessment, management, and prevention of PAD for adult patients.<sup>8 9</sup> The PAD guidelines focus on the importance of using validated tools to monitor PAD in patients. Interventions to improve the management of PAD in ICUs are associated with improvements in patient outcomes. The implementation of bundles such as the awakening and breathing coordination, delirium prevention and monitoring, and early mobility and exercise (ABCDE) saw a 3-day decrease (24 vs 21 days, p=0.04) in mechanical ventilation status,

decrease in delirium (OR=0.55, p=0.04) and increase in odds of mobilising out of beds in academic medical or surgical ICUs (OR=2.11, p=0.003).<sup>10 11</sup> Similarly, the implementation of educational and clinical interventions for elements of PAD bundles involving the interdisciplinary team improved the accuracy of PAD assessments through standardised patient assessments and improved adherence to clinical practice guidelines in an academic medical ICU.<sup>12</sup> Additionally, the introduction of online education modules resulted in a clinically relevant improvement in patient safety and was valued by nurses treating critically ill, mechanically ventilated patients.<sup>13</sup>

Despite their recognised importance, the implementation and compliance to PAD guidelines is poor, especially in community hospitals.<sup>14</sup> To assess our centre's adherence, we developed a phase I study on PAD, which engaged nurses from our community centre's adult medical-surgical ICU in a survey and focus groups that identified a care gap in PAD assessment in critically ill adult patients. The survey revealed that only 88%, 85% and 41% of nurses were comfortable with assessing PAD, respectively.<sup>15</sup> Additionally, only 47% and 42% of nurses were satisfied or strongly satisfied with PAD management in general provided by other nurses and by intensivists, respectively.<sup>15</sup> Qualitative data from this group showed divergent opinions on optimal sedation level of critically ill patients.<sup>16</sup>

Current evidence on implementation of clinical guidelines demonstrate that effective guideline implementation strategies often have multiple components.<sup>17</sup> Additionally, environmental characteristics, including support from peers and superiors, can affect guideline implementation.<sup>17</sup> Multiple studies have investigated interventions to improve the management of PAD.<sup>11–14</sup> These studies, however, have been largely conducted in academic settings, highlighting the gap in the understanding of the generalisability to community-based hospital settings. Using this information, we designed a multifaceted and multidisciplinary intervention involving nurses, family members and physicians in a community adult ICU to improve compliance with PAD guidelines.<sup>18</sup>

# **METHODS**

We used the quality improvement model using an uncontrolled before-and-after design. The intervention was developed over September 2017 to August 2018 using the Model for Improvement Plan–Do–Study–Act (PDSA) cycles.<sup>19</sup> The PDSA cycle is a rapid and iterative approach to deliver and modify interventions for the desired improvement.<sup>19</sup> For this project, the PDSA cycle approach was used to develop, test and refine the intervention components before wide scale implementation. A PAD Advisory Committee (PADAC) was formed for the development of the intervention. The PADAC included the ICU manager, five nurses, two physicians (intensivists), an ICU pharmacist and the ICU research coordinator. Patients and the public were not involved in the

BMJ Open Qual: first published as 10.1136/bmjoq-2020-001305 on 9 December 2021. Downloaded from http://bmjopenquality.bmj.com/ on April 28, 2024 by guest. Protected by copyright.

design of the study nor the choice of outcome measures. Although the public and patients were not involved in the PADAC, family members and patients provided feedback in the development of some components of our intervention. We followed the recommendations set by the Standards for Quality Improvement Reporting Excellence V.2.0 guidelines.<sup>20</sup> The protocol of this quality improvement study was published.<sup>18</sup> The goal of this study is to improve compliance to PAD assessment guidelines among nurses using a multifaceted and multidisciplinary intervention on the assessment of PAD by ICU staff in a Canadian community hospital.

This study was conducted in the ICU of St. Catharines Site, Niagara Health, a community hospital in a mediumsized city in Ontario, Canada. The site has one closed level III medical-surgical ICU with 14 beds. Approximately 10 physicians and 100 registered nurses provide ICU care. The ICU setting is described further in the study protocol.<sup>18</sup> The intervention targeted all nursing staff, family members and physicians in the ICU. The duration of the intervention was 4weeks. The nursefocused components included four educational modules developed using the 2013 PAD guidelines and visual reminders in the form of cue cards from validated assessment tools including the Critical Care Pain Observation Tool (CPOT) for pain, the Richmond Agitation-Sedation Scale (RASS) for agitation and Confusion Assessment Method for the ICU (CAM-ICU) for delirium.<sup>8</sup> The family member-focused components involved educational pamphlet regarding delirium, educational video on a dedicated computer on delirium and volunteer conducted in-person interviews of the newly admitted patients' family members. The physician-focused component included a script to remind intensivists to encourage assessment and treatment of PAD during rounds. The multidisciplinary-focused component included a poster to remind nursing staff, physicians and family members on the assessment of delirium. Descriptions of the multifaceted and multidisciplinary intervention are outlined in table 1. The specific components where possible are included in the online supplemental materials. A logic model of the multifaceted and multidisciplinary intervention is shown in online supplemental materials 6.

## Eligibility criteria and sample size calculation

All adult patients defined as ages 18 and above admitted to the ICU for more than 24 hours were included in the study. Based on a previously conducted nurse-focused quality improvement study, improvements in the proportion of patient-days with PAD assessment in pain by 8.2%, agitation by 14.4% and delirium by 14.8% are expected.<sup>21</sup> The sample size was calculated using 95% CI and power of 80% (Z beta=0.20). The minimum sample size of 277 unique patients was required.

## Study of the intervention

The effects of the PAD intervention were studied using an uncontrolled before-and-after design. The outcomes of

Table 1         Description of multifaceted and multidisciplinary intervention for pain, agitation, and delirium					
Component	Description	Refer to			
Nurse-focused components					
Educational modules	<ol> <li>4 online educational modules:</li> <li>PAD research programme, post-intensive care syndrome, basic pharmacology, pharmacokinetic properties of common ICU drugs</li> <li>Pain module: CPOT</li> <li>Agitation: RASS</li> <li>Delirium: CAM-ICU</li> <li>Followed up with weekly email reminders and demonstration of modules</li> </ol>				
Visual reminders	Validated tools including the CPOT, RASS and CAM-ICU cue cards by bedside	Online supplemental materials 1			
Family-focused components					
Interviews	Volunteer (undergraduate students) conducted in-person interviews with family members of newly admitted patients within 48–72 hours of admission Intended to empower family members to participate in PAD care by providing information on baseline cognitive function, mobility and use of visual and hearing aids	Online supplemental materials 2			
Educational pamphlet	Provided in waiting rooms to educate family members regarding delirium	Online supplemental materials 3			
Educational video	Video on dedicated computer on delirium (licensed from Osmosis.org)				
Physician-focused component					
Multidisciplinary round script	Script to remind intensivists to order target RASS Score, to discuss PAD assessment and treatment and to encourage nurses to achieve adequate pain control and light sedation	Online supplemental materials 4			
Multidisciplinary-focused component					
Poster	Reminder on the unit to assess and treat PAD	Online supplemental materials 5			

CAM-ICU, Confusion Assessment Method for the ICU; CPOT, Critical Care Pain Observation Tool; ICU, intensive care unit; PAD, pain, agitation and delirium; RASS, Richmond Agitation–Sedation Scale.

interest were selected based on recommendations from the 2013 PAD guidelines. To study the effects of the intervention, preintervention data were collected daily for 6weeks prior to the start of the intervention. Following the 4-week implementation period, postintervention data were collected daily for 6weeks. Data collection was completed by a dedicated research coordinator (MCP). Data were collected using a dedicated PAD Programme daily collection form.

The primary outcomes of this study were: (1) the proportion of patient-days with pain assessment by nurses using the Numeric Pain Rating Scale (NPRS) or the CPOT at least four times per 12-hour shift, (2) the proportion of patient-days with agitation assessment by nurses using the RASS at least four times per 12-hour shift and (3) the proportion of patient-days with delirium assessment by nurses using the CAM-ICU at least once per 12-hour shift. The secondary outcomes of this study were: (1) proportion of patient-days with significant pain defined as NPRS scores $\geq$ 4, or CPOT scores $\geq$ 3, (2) proportion of patient-days with optimal sedation levels defined by RASS scores between -2 and 0, or at target RASS Score at least 50% of the time per day, (3) proportion of patient-days with

oversedation defined by RASS score of less than -2 at least 50% of the time, (4) proportion of patient-days with agitation defined by RASS Score of greater than 0 at least 50% of the time, and proportion of patient-days with positive delirium screening using the CAM-ICU.

# Analysis

Descriptive and analytical statistics was used to analyse quantitative data, and SPSS V.26.0 was used. McNemar's test was used to detect differences on dichotomous variables between preintervention and postintervention. The level of significance was set at p<0.05.

# RESULTS

The intervention was designed and developed over a 12-month period using the Model for Improvement's PDSA test cycles. The key improvement areas and specific components of the intervention are outlined in table 2. The details of each PDSA cycle are outlined in online supplemental material 7.

Overall, 430 patient-days from 69 unique patients and 406 patient-days from 64 unique patients were included

Table 2         Initial results and development of the intervention				
Component	Key improvements	PDSA cycles		
Nurse-focused components				
Educational modules	In-person nursing engagement intervention turned to development of e-modules	4		
Visual reminders	Reviewed by MD and nurses. Feedback was collected on design and content	2		
Family-focused components				
Interviews	Volunteer conducted in-person interviews with family members of newly admitted patients, content revised	10		
Educational pamphlet	Education pamphlet developed with feedback from family and with permission from www.icudelirium.org	4		
Educational video	Purchased from Osmosis.org	3		
Physician-focused component				
Multidisciplinary round script	Script to remind intensivists to order target RASS scores, to discuss PAD assessment and treatment and to encourage nurses to achieve adequate pain control and light sedation Rephrased by physicians	5		
Multidisciplinary-focused componer	nt			
Poster	Reviewed by MDs and RNs	8		

MD, Intensivists; PAD, Pain, agitation, and delirium; RASS, Richmond Agitation-Sedation Scale; RN, registered nurse.

during the preintervention and postintervention periods, respectively, for this quality improvement project. The intervention did not improve the proportion of patientdays with guideline compliance to the assessment of pain (23.4% vs 22.4%, p=0.80), agitation (42.9% vs 38.9%, p=0.28), nor delirium (35.2% vs 29.6%, p=0.10). Of patient-days with PAD assessments, there was no significant difference in patient-days with significant pain, optimal sedation, oversedation, agitation, nor delirium (table 3).

# **Barriers to implementation**

Barriers to the implementation of our quality improvement intervention occurred on many levels. This included documentation deficiencies where nurses were documenting PAD assessment in the incorrect section of patients' charts, which may have led to the undercollection of data. There was a high rate of nursing turnover at the time of the intervention, where not all of the nurses were aware of the PAD training modules. The implementation period also coincided with hospital accreditation, which had a significant time and learning burden for ICU staff. In our setting, this reflected in poor completion rates (14%–24%) of our online educational modules. Additionally, the hospital policy lacked clarity on PAD assessment frequency that conflicted with the agitation assessment recommendation from the PAD guidelines.<sup>8</sup> Personal attributes could have also been a factor as our previous nursing survey found that only 88%, 85% and 41% of nurses were either comfortable or very comfortable with the assessment of PAD, respectively, and had differing opinions on optimal sedation.<sup>15 16</sup>

# DISCUSSION

Despite the development of guidelines for the assessment and management of PAD in the ICU, compliance to these guidelines is poor in community ICU settings.<sup>18</sup> This study evaluated a multifaceted and multidisciplinary intervention on PAD assessment in a Canadian community ICU. The intervention involving nurses, physicians and patient families resulted in no significant changes in adherence to the PAD assessment guidelines. Post intervention, the proportion of patient-days with guideline adherence remains low for pain (22.4%), agitation (38.9%) and delirium (29.6%) assessments by nurses.

There is an average 17-year gap between research and clinical practice.<sup>22</sup> Barriers to implementation in healthcare arise

 Table 3
 Preintervention and postintervention proportion of patient days with significant pain, optimal sedation, oversedation, agitation and delirium

	Preintervention	Postintervention	P value
Proportion of patient-days with significant pain	7.7%	4.9%	P=0.210
Proportion of patient-days with optimal sedation	86.1%	81.1%	P=0.081
Proportion of patient-days with oversedation	13.6%	18.9%	P=0.059
Proportion of patient-days with agitation	5.3%	4.3%	P=0.618
Proportion of patient-days with significant delirium	17.5%	15.0%	P=1.000

from many levels. In 2009, Damschroder et al established the Consolidated Framework for Implementation Research (CFIR) to guide the implementation of health services, which comprises five domains: intervention, inner settings, outer settings, individuals involved and process by which implementation is accomplished.<sup>23</sup> Specific to the ICU setting, Costa et al applied the CFIR framework in a systematic review on the implementation of the ABCDE to identify barriers to ABCDE delivery and have found barriers to implementation in all but the process category.<sup>24</sup> In our study, the CFIR framework can be applied to identify barriers to the implementation of our PAD assessment intervention. Barriers in the intervention domain included the documentation deficiencies. Barriers in the inner setting domain included high nursing turnover, the coincidence with hospital accreditation, as well as lack of clarity in existing hospital policy. We also experienced barriers in the characteristics of individuals domain, where nurses had differing comfort levels in PAD assessment and differing opinions on optimal sedation.<sup>15 16</sup> Other literature has identified barriers to delirium assessment by nurses including that it is challenging to assess in intubated patients, complexity of delirium assessment tools and that it can be time-consuming.<sup>25 26</sup>

This is one of the first studies to target not only nurses and physicians, but also patients and family members in the management of PAD in a community ICU. The formation of the PADAC ensured that ICU administrator, nursing staff and physicians were onboard and had input on the intervention, which could promote the uptake of the intervention.<sup>27</sup> This study included all patients admitted to the ICU for more than 24 hours regardless of mechanical ventilation status adding to the external validity of the intervention. The uncontrolled before-andafter design represents a pragmatic approach to investigating interventions in a resource-limited setting.

Limitations of the study include the uncontrolled before-and-after design. The study's results lacked comparisons to a control group, which implies that the intervention cannot be compared with practice changes that occur in ICU settings at baseline. The analysis of our results was completed using patient-days, and we were unable to reach our sample size due to the short study period. Due to resource limitations in community ICU settings, information on patients including baseline risk factors associated with delirium (eg, demographics, severity of illness, age, history of dementia, history of hypertension, history of delirium and substance use) was not collected to adjust for potential confounders.<sup>8</sup> Data collection was limited to patient flow sheets, while some nurses documented PAD in narrative interdisciplinary notes and were missed in data collection. The doses of medications such as sedatives, neuromuscular blockers, analgesics and antipsychotic medications were not recorded in this study. Patients were also not stratified based on mechanical ventilation status. However, our primary outcomes, that is, proportion of patient-days with guideline-recommended PAD assessments by nurses, are not influenced by the above. Other process measures

including exposure to the QI interventions that could have elucidated the effectiveness of specific components of the multifaceted and multidisciplinary intervention were not measured, and variation in physician uptake of the intervention was not assessed. This intervention focused on the compliance to PAD assessment guidelines, but did not measure the accuracy of the assessment, adherence to treatment or prevention recommendations from PAD guidelines, nor patient-outcomes.<sup>8</sup>

The current study illustrates that the implementation of a multifaceted and multidisciplinary PAD intervention did not improve adherence to PAD assessment guidelines by nurses in a Canadian community ICU. The rate of PAD assessment remains low postintervention. PAD experienced in the ICU has important implications on patient outcomes, including patient mortality. Therefore, the reliable detection and diagnosis of PAD is essential for PAD treatment and ongoing interventions are necessary to improve guideline adherence. Components of the intervention are still in place, including educational modules for nurses to complete, as well as the multidisciplinary poster. Our quality improvement intervention identified barriers specific to our ICU that need to be addressed in future iterations. This includes inner setting, intervention, and individuals involved domains such as: (1) including the education modules in the orientation of new nurses and reintroduction of mandatory nurse education modules using organisation approved platforms to assist completion, (2) updating the ICU policy to include the frequency of PAD assessment and its inclusion of PAD assessment in ICU admission order sets, (3) including PAD assessment frequency and target RASS scores in pre-printed ICU admission order sets, (4) conducting an audit to evaluate whether continuation of intervention would improve outcomes, (5) surveying nurses on perceptions on the intervention to potentially simplifying the intervention to increase buy-in and (6) carrying the study on to the quality improvement division of the hospital to improve buy-in. Future implementation of PAD guidelines should assess their own units' specific barriers to PAD implementation and design specific interventions tailored to overcome these barriers with consideration of the context, team and capacity for implementation.

### Author affiliations

<sup>1</sup>Niagara Regional Campus, McMaster University Michael G DeGroote School of Medicine, St. Catharines, Ontario, Canada

<sup>2</sup>Niagara Health System—Saint Catharines Site, Saint Catharines, Ontario, Canada <sup>3</sup>Department of Health Science, Brock University, Saint Catharines, Ontario, Canada <sup>4</sup>Department of Health Research Methods, Evidence, and Impact, McMaster University, Hamilton, Ontario, Canada

<sup>5</sup>Medicine/Critical Care, McMaster University Department of Medicine, Hamilton, Ontario, Canada

<sup>6</sup>Medicine/Critical Care, Niagara Health, St. Catharines, Ontario, Canada Twitter Madelyn Law @MadelynLaw

Acknowledgements JLYT is the recipient of the McMaster University, Department of Medicine Mid-Career Research Award.

**Contributors** ZM analysed the data, drafted and revised the manuscript. MC implemented the intervention including modifying the intervention based on feedback from staff. MC collected data from the unit. ML contributed to the

# **Open** access

intervention design, implementation using quality improvement methodology and revisions of the manuscript. DJ contributed to the statistical design and analysis of the results. H-OC contributed to the implementation of the intervention and revision of the manuscript. JNHL contributed to the revision of the manuscript. JLYT is the guarantor of the project and contributed to the conception and design of the study, implementation of the interventions, and writing and revisions of the manuscript.

Funding Funding was provided by the Physician Services Incorporated Community Research Fund and the Medbuy Research and Education Fund. No award/grant number.

Competing interests None declared.

#### Patient consent for publication Not required.

Ethics approval This study involves human participants and was approved by The Hamilton Integrated Research Ethics Board approved this study (18-040C). The study was approved by the Hamilton Integrated Research Ethics Board (18-040-C), with a waiver on the need to obtain consent as this is a quality improvement study. All identifying patient information was anonymizsed and stored in password-protected computers in a secured research office.

Provenance and peer review Not commissioned; externally peer reviewed.

Data are available upon request.

Supplemental material This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

**Open access** This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: http://creativecommons.org/licenses/by-nc/4.0/.

### **ORCID iDs**

Zechen Ma http://orcid.org/0000-0003-3064-8118 Jennifer LY Tsang http://orcid.org/0000-0003-1935-0791

## REFERENCES

- Erstad BL, Puntillo K, Gilbert HC, et al. Pain management principles in the critically ill. Chest 2009;135:1075–86.
- 2 Chanques G, Sebbane M, Barbotte E, et al. A prospective study of pain at rest: incidence and characteristics of an unrecognized symptom in surgical and trauma versus medical intensive care unit patients. *Anesthesiology* 2007;107:858–60.
- Siegel MD. Management of agitation in the intensive care unit. *Clin Chest Med* 2003;24:713–25.
- 4 Salluh JIF, Wang H, Schneider EB, et al. Outcome of delirium in critically ill patients: systematic review and meta-analysis. BMJ 2015;350:h2538–1833.
- 5 Ely EW, Shintani A, Truman B, et al. Delirium as a predictor of mortality in mechanically ventilated patients in the intensive care unit. JAMA 2004;291:1753–62.
- 6 McNicoll L, Pisani MA, Zhang Y, *et al*. Delirium in the intensive care unit: occurrence and clinical course in older patients. *J Am Geriatr* Soc 2003;51:591–8.
- 7 Pisani MÁ, Kong SYJ, Kasl SV, *et al.* Days of delirium are associated with 1-year mortality in an older intensive care unit population. *Am J Respir Crit Care Med* 2009;180:1092–7.
- 8 Barr J, Fraser GL, Puntillo K, et al. Clinical practice guidelines for the management of pain, agitation, and delirium in adult patients in the intensive care unit. Crit Care Med 2013;41:263–306.

- 9 Devlin JW, Skrobik Y, Gélinas C, et al. Clinical practice guidelines for the prevention and management of pain, Agitation/Sedation, delirium, immobility, and sleep disruption in adult patients in the ICU. *Crit Care Med* 2018;46:e825–73.
- 10 Balas MC, Burke WJ, Gannon D, et al. Implementing the awakening and breathing coordination, delirium monitoring/management, and early exercise/mobility bundle into everyday care: opportunities, challenges, and lessons learned for implementing the ICU pain, agitation, and delirium guidelines. Crit Care Med 2013;41:S116–27.
- 11 Balas MC, Vasilevskis EE, Olsen KM, et al. Effectiveness and safety of the awakening and breathing coordination, delirium monitoring/ management, and early exercise/mobility bundle. Crit Care Med 2014;42:1024–36.
- 12 Weber ML, Byrd C, Cape K. Implementation of the ABCDEF bundle in an academic medical center. *Journal of Clinical Outcomes Management* 2017;24.
- 13 Walsh TS, Kydonaki K, Antonelli J, et al. Staff education, regular sedation and analgesia quality feedback, and a sedation monitoring technology for improving sedation and analgesia quality for critically ill, mechanically ventilated patients: a cluster randomised trial. *Lancet Respir Med* 2016;4:807–17.
- 14 Yan W, Morgan BT, Berry P, et al. A quality improvement project to increase adherence to a pain, agitation, and delirium protocol in the intensive care unit. *Dimens Crit Care Nurs* 2019;38:174–81.
- 15 Maximous R, Miller F, Tan C, et al. Pain, agitation and delirium assessment and management in a community medical-surgical ICU: results from a prospective observational study and nurse survey. BMJ Open Qual 2018;7:e000413.
- 16 Tsang JLY, Ross K, Miller F, et al. Qualitative descriptive study to explore nurses' perceptions and experience on pain, agitation and delirium management in a community intensive care unit. BMJ Open 2019;9:e024328.
- 17 Francke AL, Smit MC, de Veer AJE, et al. Factors influencing the implementation of clinical guidelines for health care professionals: a systematic meta-review. BMC Med Inform Decis Mak 2008;8:38.
- 18 Penuela MC, Law M, Chung H-O, et al. Impact of a multifaceted and multidisciplinary intervention on pain, agitation and delirium management in a Canadian community intensive care unit: a quality improvement study protocol. CMAJ Open 2019;7:E430–4.
- 19 Langley G, Moen Ř, Nolan K. *The improvement guide: a practical approach to enhancing organizational performance*. 2nd ed. San Francisco: Jossey-Bass Publishers, 2009.
- 20 Ogrinc G, Davies L, Goodman D, et al. Squire 2.0 (standards for quality improvement reporting excellence): revised publication guidelines from a detailed consensus process. *Perm J* 2015;19:65–70.
- 21 Tan CM, Camargo M, Miller F, et al. Impact of a nurse engagement intervention on pain, agitation and delirium assessment in a community intensive care unit. BMJ Open Qual 2019;8:e000421.
- 22 Morris ZS, Wooding S, Grant J. The answer is 17 years, what is the question: understanding time lags in translational research. *J R Soc Med* 2011;104:510–20.
- 23 Damschroder LJ, Aron DC, Keith RE, et al. Fostering implementation of health services research findings into practice: a consolidated framework for advancing implementation science. *Implement Sci* 2009;4:50.
- 24 Costa DK, White MR, Ginier E, et al. Identifying barriers to delivering the awakening and breathing coordination, delirium, and early Exercise/Mobility bundle to minimize adverse outcomes for mechanically ventilated patients: a systematic review. *Chest* 2017;152:304–11.
- 25 Devlin JW, Fong JJ, Howard EP, et al. Assessment of delirium in the intensive care unit: nursing practices and perceptions. Am J Crit Care 2008;17:555–65.
- 26 Guttormson JL, Chlan L, Tracy MF, *et al*. Nurses' attitudes and practices related to sedation: a national survey. *Am J Crit Care* 2019;28:255–63.
- 27 Barr J, Pandharipande PP. The pain, agitation, and delirium care bundle: synergistic benefits of implementing the 2013 pain, agitation, and delirium guidelines in an integrated and interdisciplinary fashion. *Crit Care Med* 2013;41:S99–115.