

BMJ Open Quality Decreasing driveline infections in patients supported on ventricular assist devices: a care pathway approach

Julia Seretny,¹ Tara Pidborochynski,² Holger Buchholz,³ Darren H Freed,^{3,4} Roderick MacArthur,³ Nicole Dubyk,⁵ Laura Cunliffe,⁵ Osiris Zelaya,⁵ Jennifer Conway^{2,6}

To cite: Seretny J, Pidborochynski T, Buchholz H, *et al*. Decreasing driveline infections in patients supported on ventricular assist devices: a care pathway approach. *BMJ Open Quality* 2022;**11**:e001815. doi:10.1136/bmjopen-2022-001815

► Additional supplemental material is published online only. To view, please visit the journal online (<http://dx.doi.org/10.1136/bmjopen-2022-001815>).

Received 11 January 2022
Accepted 17 May 2022



© Author(s) (or their employer(s)) 2022. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

¹Faculty of Nursing, University of Alberta, Edmonton, Alberta, Canada

²Pediatric Cardiology, University of Alberta Faculty of Medicine & Dentistry, Edmonton, Alberta, Canada

³Department of Surgery, University of Alberta Faculty of Medicine & Dentistry, Edmonton, Alberta, Canada

⁴Division of Pediatric Cardiac Surgery, Stollery Children's Hospital, Edmonton, Alberta, Canada

⁵Mazankowski Alberta Heart Institute, Edmonton, Alberta, Canada

⁶Division of Pediatric Cardiology, Stollery Children's Hospital, Edmonton, Alberta, Canada

Correspondence to

Dr Jennifer Conway;
jennifer.conway2@ahs.ca

ABSTRACT

Background Driveline infections (DLIs) are a common adverse event in patients on ventricular assist devices (VADs) with incidence ranging from 14% to 59%. DLIs have an impact on patients and the healthcare system with efforts to prevent DLIs being essential. Prior to our intervention, our program had no standard driveline management presurgery and postsurgery. The purpose of this Quality Improvement (QI) initiative was to reduce DLIs and related admissions among patients with VAD within the first year post implant.

Methods In anticipation of the QI project, we undertook a review of the programs' current driveline management procedures and completed a survey with patients with VAD to identify current barriers to proper driveline management. Retrospective data were collected for a pre-QI intervention baseline comparison group, which included adult patients implanted with a durable VAD between 1 January 2017 and 31 July 2018. A three-pronged care pathway (CP) was initiated among patients implanted during August 2018 to July 2019. The CP included standardised intraoperative, postoperative and predischarge teaching initiatives and tracking. Using statistical process control methods, DLIs and readmissions in the first year post implant were compared between patients in the CP group and non-CP patients. P-charts were used to detect special cause variation.

Results A higher proportion of CP group patients developed a DLI in the first year after implant (52% vs 32%). None developed a DLI during the index admission, which differed from the non-CP group and met criteria for special cause variation. There was a downward trend in cumulative DLI-related readmissions among CP group patients (55% vs 67%). There was no association between CP compliance and development of DLIs within 1 year post implant.

Conclusion The CP did not lead to a reduction in the incidence of DLIs but there was a decrease in the proportion of patients with DLIs during their index admission and those readmitted for DLIs within 1 year post implant. This suggests that the CP played a role in decreasing the impact of DLIs in this patient population. However, given the short time period of follow-up longer follow-up will be required to look for sustained effects.

INTRODUCTION

In North America, diseases of the heart and heart failure are one of the leading causes of death.¹ Transplantation is the gold standard of

WHAT IS ALREADY KNOWN ON THIS TOPIC?

⇒ Driveline infections (DLIs) in durable patients with ventricular assist device (VAD) are costly adverse events for the healthcare system and the patient. Despite this, to date, only one recently published protocol on driveline management exists.

WHAT THIS STUDY ADDS?

⇒ Targeted Quality Improvement initiatives can help to reduce the burden of DLIs among patients supported on durable VADs.

HOW THIS STUDY MIGHT AFFECT RESEARCH, PRACTICE OR POLICY?

⇒ Management of the driveline requires a multidisciplinary effort that begins intraoperatively and continues postoperatively with ongoing patient management and education.

care for patients with advanced heart failure. However, the availability of donor organs is not sufficient to meet demand and certain comorbidities preclude some patients from transplant candidacy. To circumvent this, ventricular assist devices (VADs) can serve as a bridge to transplantation (BTT) or as destination therapy in this patient population.^{2–5} Yet, VAD support is not without its complications, with driveline infections (DLIs) being among the most commonly reported adverse event.^{6–11}

Various studies have found that the incidence of DLIs in patients with VAD ranges from 14% to 59%, while other studies have reported that up to 60% of unplanned hospital readmissions among patients with VAD are due to DLIs.^{6 7 9 10 12–16} Risk factors associated with development of DLIs have been highly variable and at times inconsistent. Some studies have reported that obesity exists as a predictor of DLIs^{17 18} while other studies have not replicated this finding.¹⁹ Other reported risk factors include younger age,¹³ older age,²⁰ diabetes,²⁰ depression,⁸

renal failure⁸ and driveline and device characteristics.^{21–23} DLIs often occur early after implantation suggesting that the index hospital admission may play an important role in the development of infections.²⁴ When aiming to reduce or eliminate the risk of DLI in patients with VAD, the variability in identified risk factors serves as a challenge for singular interventions.

DLIs decrease quality of life as they are a significant source of morbidity and mortality.^{6, 7, 9} Furthermore, hospital readmissions due to DLIs negatively affect the healthcare system as they require intensive management, diagnostic testing, further medical interventions and increase healthcare costs.^{14, 15, 25} A recent study identified the median direct hospital cost corresponding with a DLI-related readmission as 11 506 US\$.²⁶ Not surprisingly, DLIs are an expensive and undesired outcome not only for patients and care teams, but also for the healthcare system itself.²⁶

Guidelines for the care of drivelines has been lacking with the first published protocol coming out in 2020.²⁷ Our programme had no comprehensive standard approach to driveline care and over a 5-year period (2013–2018), there were 278 readmissions among 79 patients with VAD. Of these readmissions, 51% (n=141) were for VAD-related reasons, with the most common being DLIs (37%) with 52 DLI-related readmissions recorded during this time period. The median length of stay (LOS) for a DLI was 9 days (IQR 7 to 12) with a range from 2 to 88 days. Based on the average daily cost of a hospital admission during this time period and using the median duration of a DLI-related readmission LOS, we inferred that DLI readmissions cost our healthcare system ~\$1 million to \$3.4 million (CDN) depending on the location of readmission and stay (ie, ward vs cardiovascular intensive care unit (CVICU)).

As the causes of DLIs are multifactorial, we aimed to address this issue through a collaborative and preventative approach. We carried out a Quality Improvement (QI) project that sought to develop and implement a standardised three-pronged care pathway (CP) with the goal of reducing DLI-related readmissions in the first year post-VAD implant.

METHODS

Setting

The adult Artificial Heart Programme at the Mazankowski Alberta Heart Institute, University of Alberta Hospital (Edmonton, Alberta, Canada) has been implanting durable VADs with a driveline since 2009 and serves as a referral centre for other provinces in western Canada. From June 2009 through July 2019, 164 patients received a durable VAD that had a driveline. The programme consists of a multidisciplinary team of stakeholders: cardiac surgeons, anaesthetists, VAD coordinators, CVICU and ward nurses, CVICU intensivists, social workers, dietitians, as well as patient and family caregivers.

Study design

Our QI project focused on reducing DLI infections and related admissions among adult patients on durable VAD support. Formulation of the QI project began in January 2018. In anticipation of the project, we undertook several preintervention analyses: (1) a review of the programme's current driveline management procedures; (2) a survey with patients with VAD to identify barriers to proper driveline management and (3) a retrospective chart review to determine the baseline incidence of DLIs in patients implanted with a VAD. Based on the data from the preintervention analyses, we developed a three-pronged CP that was practice-based, with initiatives and education centred on the medical management of the driveline following VAD implantation. The QI project included adult patients undergoing VAD implantation between 1 August 2018 and 31 July 2019 (CP group) with patients followed for 1 year after implantation. At the end of the QI project, we conducted chart abstraction of patients involved in the CP to determine the incidence of DLIs and related hospital readmissions. Results from the CP group were compared with the retrospective cohort (non-CP group) to determine if the CP led to an overall reduction in incidence of DLIs and related hospital readmissions.

Preintervention review of driveline management procedures

Prior to the initiation of this QI project, there was no standard intraoperative procedure for driveline securing or velour positioning and there was variability in antibiotic prophylaxis at the time of VAD implant. Postoperatively, daily dressing changes occurred for the first 10 days followed then by dressing changes Monday/Wednesday/Friday dressing with patients and family receiving didactic teaching about driveline care and being observed performing the dressing change before patient discharge.

Preintervention patient survey

A convenience sample of patients was surveyed to identify common barriers that patients and families experience with caring for the driveline. Seven patients provided answers to the four-question survey and the consistent theme identified was that the cost of the dressing kit and supplies was a major barrier to proper driveline dressing management, with some patients indicating that they used alternatives to the recommended supplies in order to reduce the cost. Based on this feedback, the VAD programme developed a new, more cost effective driveline dressing kit. The cost of the new kit was \$13CDN compared with the previous cost of \$30CDN per kit.

Preintervention retrospective chart review

To serve as a baseline comparison, retrospective data were collected for a pre-CP intervention group, which included adult patients implanted with a durable VAD between 1 January 2017 and 31 July 2018 (non-CP group).

Table 1 CP Initiatives and Compliance Scoring Metric

| Time period | CP Standardised Initiative | Compliance Score points |
|----------------|--|-------------------------|
| Intraoperative | Use of dual antibiotics (vancomycin and cefazolin) at the time of VAD implant | 1 |
| | Velour positioning 1 cm from exit site | 1 |
| | Two internal sutures used to secure the driveline (4–0 Vicryl); two external staples for skin closure at driveline exit site | 1 |
| | Use of dual antibiotics (vancomycin and cefazolin) postoperatively | 1 |
| Postoperative | After initial driveline dressing change (48 hours post implant), dressing changes to occur Monday, Wednesday and Friday | 1 |
| | Delay in showering for 10 days postoperatively | 1 |
| | Dietitian counselling with the patient about obesity, weight gain and risk of DLI occurrence | 1 |
| | Social work counselling about patient-specific barriers to driveline dressing changes | 1 |
| Predischarge | Comprehensive patient/caregiver teaching of driveline management including: <ul style="list-style-type: none"> ▶ Routine care of driveline ▶ Proper positioning to prevent pull at the exit site ▶ Tips to prevent DLIs ▶ Signs and symptoms of DLIs ▶ How to notify VAD Coordinators if signs/symptoms are present | 2 |
| | Driveline dressing brochure and infographic pamphlets given to the patient/caregiver | 1 |
| | Driveline dressing video watched at two separate time-points | 2 |
| | Driveline dressing done by patient/caregiver at two separate time-points | 2 |
| | Driveline dressing quiz completed at two separate time-points | 2 |
| | Total score | |

CP, care pathway; DLI, driveline infection; VAD, ventricular assist device.

QI care pathway interventions

The QI technique used was a Total Quality Management Approach where emphasis is on a team-based approach for quality improvement.²⁸ A DLI QI team was created that included VAD physicians, surgeons, coordinators, a social worker and a dietician who worked towards standardising driveline care and management.

As one of our primary drivers of change in this QI project, our goal was to develop and implement a three-pronged CP that was practice-based, with initiatives and education centred on the medical management of the driveline following VAD implantation.

Initiatives within the CP targeted cardiac surgeons, multidisciplinary team members (eg, dietitians, social workers, nurses), as well as patients and caregivers. The focus of the CP was on standardising driveline management at three time-points: intraoperative, postoperative and predischarge. We developed the CP after a number of meetings among the DLI QI team, review of the literature^{6 29–37} and feedback from stakeholders. Prior to implementation of the CP, the QI initiative was presented to stakeholders involved in the project in order to orient them to the initiative and outline their roles.

Table 1 provides an overview of all time-points and initiatives included as part of the CP.

One aspect of the CP was to foster education and awareness among the medical team and patients/caregivers

about driveline management and infections. As a part of this, we developed a Driveline Dressing Video to use as an accessible educational resource. This video provided detailed footage of a driveline dressing change occurring from start to finish following the standardised protocol and aseptic technique. Further, we developed a digital Driveline Dressing Quiz that was to be used in conjunction with the video as an additional educational resource. The quiz was a direct reflection of the information presented in the video and was composed of 17 questions with a total score of 23 points. This quiz was tested on the VAD coordinators and other stakeholders (eg, social workers, dietitians) prior to its finalisation to assess difficulty, ease of completion and relateness to the Driveline Dressing Video. The video and quiz were uploaded to a program-specific iPad for patient and/or caregivers to review prior to completing their own driveline dressing changes in hospital and prior to discharge. In addition, new paper-based infographics for patient use were developed that visually focused on the routine care of a driveline, proper positioning of the driveline to prevent any pull at the exit site, tips to prevent DLIs, signs and symptoms of DLIs and how to notify the VAD team if any signs or symptoms of infection were present.

Targeted specifically to the medical team, CVICU as well as ward nurses attended a 15 min in-service held by the VAD coordinators to review the new dressing change

protocol and proper sterile technique for a VAD dressing change. For ease of access and as a source of ongoing reference among nursing staff, the Driveline Dressing Video was uploaded to CVICU and ward desktop workstations. Review of VAD driveline dressing changes and review of the video and quiz were included in the nurses' annual VAD re-certification.

To track compliance with the CP, standardised Driveline Checklists encompassing 17 components of the initiative were created for use with VAD in-patients (table 1). Use of the newly developed low-cost dressing kit was not included in the compliance score. Missing documentation in a patient's chart regarding completion of an initiative was interpreted as the initiative was not followed by a member of the team. Each patient received a compliance score out of 17 points, with good compliance to the CP protocol considered a score $\geq 80\%$.³⁸

DLI data collection

Data collection for both groups (preintervention group (non-CP) and CP) included patient demographics, clinical and VAD characteristics, outcome at the end of follow-up, as well as DLI-related information up to 1 year post implant. DLIs were defined as per the Intermacs definitions which included clinical symptoms accompanied by a positive culture and initiation of antibiotic treatment.^{39 40}

Measures

The outcome measures for the study included the incidence of first DLI, incidence of DLIs that occurred during the patient's index admission and occurrence of DLI-related hospital readmissions within 1 year post implant. Our process measure included the overall compliance score as based on adherence to the intraoperative, postoperative and predischarge initiatives and standardised protocols by clinical professionals, patients and caregivers.

Data analysis

SPSS V.27.0 (IBM, USA) was used for descriptive statistical comparisons between the non-CP and CP groups. Descriptions for continuous variables are reported in terms of proportion, median, and the 25th, 75th IQR. To assess differences between study groups, the Mann-Whitney U (Wilcoxon rank-sum) test (two-tailed p value) was used to assess continuous variables whereas the Fisher's exact test (one-tailed p value) was used to assess differences between categorical variables.

Time series data were analysed using statistical process control methods using P-Charts and run charts. These types of charts were used to compare DLIs and readmissions over time in the first year post implant between the non-CP group and CP group. Specifically, P-Charts, graphing proportion values over time, were used to detect special cause variation attributable to the implementation of the CP. Benneyan *et al* define special cause variation as 'unnatural variation due to events, changes

or circumstances that have not previously been typical or inherent in the regular process' (p.59).⁴¹ This type of variation, as opposed to common cause variation, would indicate if the variations in our outcome measures were attributable to the implementation of the CP or occurred inherently and predictably.^{41 42} P-charts were analysed using standard criteria for probability established for assessing data points and their behaviours.⁴¹ The upper and lower limits were set at three SD from the mean. Run charts were used to examine variation over time for compliance scores with the median score determined and plotted as the centerline and probability rules used to determine special cause variation.

Patient and public involvement

The development of the research question and quality improvement project was designed to reduce the occurrence of DLIs and related hospital admissions in patients implanted with durable VADs. Patients and the public were not involved in any aspect of the research study (eg, design, recruitment, conduct). Dissemination of results is intended through open access publication.

RESULTS

Demographic and clinical characteristics

A total of 55 patients were included in the project (non-CP (n=34) and CP (n=21) group). Online supplemental table 1 shows baseline, clinical and DLI characteristics for all patients with a comparison of the non-CP and CP groups separately. The median age was 56 years (IQR 45 to 61) with the majority (87%) of the cohort male. The majority (87%) of patients had a diagnosis of cardiomyopathy, with idiopathic cardiomyopathy identified as the most common (45%) aetiology. Among all patients, the most common therapeutic intent for VAD use was as a BTT (86%). Overall, most patients (78%) were implanted with the HeartMate III (HM3) (Abbott, Abbott Park, Illinois, USA) and the median duration of support was 601 days (IQR 405 to 825). There was no significant difference in demographics and clinical characteristics between the two groups (online supplemental table 1). Just under half (47%) of the patients were still on VAD support when the last patient in the CP group reached the 1-year follow-up time-point.

DLIs during follow-up period

As shown in online supplemental table 1, a higher proportion of patients in the CP versus non-CP group developed a DLI within the first year post-VAD implant (52% vs 32%, $p=0.12$). The median time to first DLI post implant was longer in the CP group (147 days; IQR 97 to 280) than in the non-CP group (88 days; IQR 53 to 202), although not statically significant ($p=0.19$) (online supplemental table 1). In addition, a smaller proportion of patients in the CP group (55%) were readmitted to hospital within the first year post implant with a DLI than those in the non-CP group (67%) ($p=0.47$).




Figure 1 Control chart displaying the cumulative proportion of implanted patients with a DLI, with n representing the cumulative number of implanted patients. Only first DLIs were included. The central red line represents the mean with upper and lower limits set at 3 SDs from the mean (dashed lines). CP, care pathway; DLI, driveline infection; VAD, ventricular assist device.

When examined using statistical process control methods, the P-Chart in [figure 1](#) highlights an increasing trend for cumulative DLIs post-CP implementation. However, no DLIs developed during the index hospital stay among patients in the CP group ([figure 2](#)), which differed from the non-CP group and met criteria for special cause variation. Lastly, a downward shift in the cumulative DLI-related readmissions among patients in the CP group was noted ([figure 3](#)) but this shift did not meet the criteria for special cause variation.

Protocol compliance

When analysing compliance scores, 62% of the patients had a compliance score of $\geq 80\%$. The median compliance score was 65% (IQR 27 to 94) ([figure 4](#)). No association was seen between good compliance (score $\geq 80\%$) and development of a DLI within 1 year post implant ($p=0.60$) or readmission ($p=0.25$). As [figure 4](#) highlights, there was a noticeable decrease in compliance over time, with the last two quarters of the project having the worst




Figure 2 Control chart displaying the cumulative proportion of DLIs that occurred during index admission, with n representing the number of DLIs that occurred by the end of the quarter (Q). The central red line represents the mean with upper and lower limits set at 3 SDs from the mean (dashed lines). CP, care pathway; DLI, driveline infection; VAD, ventricular assist device.




Figure 3 Control chart displaying the cumulative proportion of DLIs that required Hospital readmission, with n representing the number of readmissions in the quarter (Q). Each patient was censored following their first admission to hospital (excludes those DLIs occurring during index admission). CP, care pathway; DLI, driveline infection; VAD, ventricular assist device.

compliance scores and the majority of DLIs in the CP group (66%). The shift in compliance scores below the median almost satisfies the criteria for special cause variation, with the exception of one outlier in Quarter 3 of 2019.

DISCUSSION

This QI project standardised driveline management for patients with VAD at three interdisciplinary time-points, in addition to initiating standard and enhanced patient and provider education, as well as documentation tools

within this programme. Although there was an increase in DLIs among patients involved with the CP, there was a reduction in DLI-related readmissions over the first year post implant. Furthermore, there also was a significant decrease in DLIs during the index admission among patients in the CP group. Thus, indicating that the implementation of the CP did in fact play a role in reducing the impact of DLIs in this patient population within our programme. Additionally, the increased time to infection may suggest that the multiprong interdisciplinary initiatives were effective initially in addition to




Figure 4 Run chart displaying compliance scores with the CP protocol for each patient with median displayed in blue. Red dots represent patients implanted in that quarter (Q) that developed a DLI within a year of VAD implant. CP, care pathway; DLI, driveline infection; VAD, ventricular assist device.

standard of care but that the impact diminished over time as patient management occurred in the outpatient setting. Lastly, it is possible that the increased incidence of DLIs among patients in the CP group was impacted by the change in implanted device during the duration of the QI project. Prior to the initiation of the CP, the predominant devices implanted were the HeartWare VAD (HVAD) (Medtronic, Minneapolis, Minnesota, USA) and the HeartMate II (HMII) (Abbott, Chicago, Illinois, USA) but around the time the CP was started, the programme had switched to implanting patients predominantly with the HM3. Compared with the HVAD and HMII, the HM3 has a driveline that is larger in diameter and is stiffer which may increase the risk of patients implanted with this type of device developing DLIs.^{22 27 43} This is relevant to our study given that all the patients in the CP group were implanted with the HM3 device and we saw an increased incidence of DLIs among patients in the CP group compared with the non-CP group where only 65% of patient were implanted with the HM3.

The improvements in terms of reducing DLIs during the index admission and DLI-related readmissions within the first year post implant among CP patients illustrate the effectiveness of addressing driveline management through an interdisciplinary approach. This aligns with a study conducted by Bernhardt *et al* in which a 10-step standard operating procedure (SOP) for driveline exit site (DLES) care (the DESTINE SOP) and wound staging was cultivated.²⁷ Similar to our study, key aspects included standardising procedures for dressing changes, driveline immobilisation, patient training and re-evaluation. We agree with Bernhardt *et al* that an interdisciplinary approach to DLES management and awareness is needed to prevent DLIs as well as increase patient compliance and quality of life.²⁷ While the above article had some similarities with respect to approach to driveline management, the study did not evaluate the impact of their QI initiative.

Another key finding was that compliance scores decreased over time and ultimately the worst scores coincided with the period in which the majority of DLIs in the CP group occurred. This phenomenon, where continued implementation of an initiative waivers over time, has been well described and is sometimes referred to as the 'bathtub curve', where the impact may decrease overtime for design or situational factors.³⁸ In this case, situational factors such as changing team members and burnout over time may have played a role. Successful CP uptake and implementation is a complex process and requires careful consideration about facilitators and barriers in order to change behaviour.⁴⁴ Facilitators and barriers to the uptake of our CP need to be further explored. Our patterns of compliance may indicate that further resources are needed at adapting implementation strategies over time based on feedback from CP stakeholders.

By taking a QI approach, this allowed us to not only assess the impact of DLIs and readmissions in the patient population but also actively implement

initiatives intended to target aspects of care in the hopes of improving short-term and long-term outcomes. The QI approach required consideration of all stakeholders and interdisciplinary team members as our initiatives may directly affect them. Instead of simply collecting data on VAD-related readmissions, we dove deeper to assess how we could effectively improve our target population's experience with DLIs and readmissions. Compared with traditional statistical methods where a measure of central tendency is compared before and after an intervention, QI methods look for trends over time, which may be missed by statistical tests. Run and control charts help us better understand the nature of the data and look for trends of improvement or worsening with interventions and if the impact is sustainable.^{4 45 46} This is clearly highlighted in this QI initiative where traditional statistical methods did not show any significant difference in outcomes pre-intervention and post interventions. However, by understanding the variance over time we were able to observe changes through the QI methods employed.

Limitations

There are a number of limitations inherent within the study design that may have affected the outcomes. CP development and implementation involves the utilisation of significant resources.⁴³ Following completion of the first cycle, we began to focus on ways to improve the CP. However, despite some modifications to the CP, we were unable to proceed with a second cycle given the beginning of the COVID-19 pandemic, which resulted in decreased resources for the programme, a decrease in surgeries and redeployment of key staff. We felt that all these factors would likely have a major impact on the overall result of a second cycle.

Given the time span of the QI project, it was impossible to control for the change in VAD device used by our centre. With this, there is potentially an inherent selection bias that could not have been predicted at the start of this project as there was little information available about the HM3 and it was not known that this device would become the primary device implanted at our institution. There are limitations to the generalisability of the outcomes found in our study due to the small sample size. The small sample size also resulted in low statistical power, which not only limited inferential statistical assessment but also likely affected our results from the statistical process control methods, as there were less than the number recommended of data points post intervention. In addition, there were 12 patients that did not meet the 1-year follow-up point due to explant of the device, transplant or death and therefore may have resulted in bias to the sample. Lastly, the CP involved multiple components and it is not possible to tease out which components were the most useful and which components need further refinement. However, from the data, it appears that the CP was successful at preventing in-hospital DLIs during the index admission and at reducing DLI-related admissions although there were more patients with DLIs

within the first year post implantation compared with the non-CP group. This highlights an area for the programme to focus on which is outpatient care of the driveline so that the initial improvements after surgery with respect to DLIs is sustained as the patient is followed postdischarge in the outpatient setting.

CONCLUSION

With the observed improvements, we speculate that the CP induced a cultural shift in awareness regarding the importance of effective and standardised techniques for driveline management, infection identification and prevention. This shift in awareness likely played an important role in reducing the impact of DLIs for this patient population but further work is required to maintain and improve compliance over time to ensure ongoing effectiveness.

Contributors All authors contributed to the preparation of this manuscript. Material preparation and data collection were performed by JS, ND, LC and OZ. JS wrote the first draft of the manuscript with all other authors commenting on previous versions. All authors read and approved the final manuscript.

Funding Alberta Health Service Chief Medical Officer (CMO) Quality Innovation Fund.

Competing interests HB contributed to Consultant for Abbott. JC contributed to non-restricted education grant from Abbott; Medical Monitor for Pumpkin Trial. DHF contributed to Stock Owner for Tevosol.

Patient and public involvement Patients and/or the public were not involved in the design, or conduct, or reporting, or dissemination plans of this research.

Patient consent for publication Not applicable.

Ethics approval This study involves human participants but based on the criteria established by the Research Ethics Board at the University of Alberta (Edmonton, Alberta, Canada), this project met the criteria for Quality Improvement and as such was exempt from full ethical review.

Provenance and peer review Not commissioned; externally peer reviewed.

Data availability statement Data are available upon reasonable request.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

REFERENCES

- Statistics Canada. Death and causes of death, 2014. The daily 2017; component of statistics Canada Catalogue No. 11-001-X.
- Miller LW, Rogers JG. Evolution of left ventricular assist device therapy for advanced heart failure: a review. *JAMA Cardiol* 2018;3:650–8.
- Kormos RL, Cowger J, Pagani FD, et al. The Society of thoracic surgeons Intermacs database annual report: evolving indications, outcomes, and scientific partnerships. *Ann Thorac Surg* 2019;107:341–53.
- Kirklin JK, Xie R, Cowger J, et al. Second annual report from the ISHLT mechanically assisted circulatory support registry. *J Heart Lung Transplant* 2018;37:685–91.
- Slaughter MS, Pagani FD, McGee EC, et al. HeartWare ventricular assist system for bridge to transplant: combined results of the bridge to transplant and continued access protocol trial. *J Heart Lung Transplant* 2013;32:675–83.
- Yarboro LT, Bergin JD, Kennedy JLW, et al. Technique for minimizing and treating driveline infections. *Ann Cardiothorac Surg* 2014;3:557–62.
- Hernandez GA, Breton JDN, Chaparro SV. Driveline infection in ventricular assist devices and its implication in the present era of destination therapy. *Open J Cardiovasc Surg* 2017;9:1–6.
- Gordon RJ, Weinberg AD, Pagani FD, et al. Prospective, multicenter study of ventricular assist device infections. *Circulation* 2013;127:691–702.
- Gordon RJ, Quagliarello B, Lowy FD. Ventricular assist device-related infections. *Lancet Infect Dis* 2006;6:426–37.
- Cagliostro B, Levin AP, Fried J, et al. Continuous-Flow left ventricular assist devices and usefulness of a standardized strategy to reduce drive-line infections. *J Heart Lung Transplant* 2016;35:108–14.
- Krzeli K, Petricevic M, Gasparovic H. Ventricular assist device driveline infections: a systematic review. *Thorac Cardiovasc Surg* 2021.
- Leuck A-M. Left ventricular assist device driveline infections: recent advances and future goals. *J Thorac Dis* 2015;7:2151–7.
- Goldstein DJ, Naftel D, Holman W, et al. Continuous-Flow devices and percutaneous site infections: clinical outcomes. *J Heart Lung Transplant* 2012;31:1151–7.
- Smedira NG, Hoercher KJ, Lima B, et al. Unplanned Hospital readmissions after HeartMate II implantation. *JACC: Heart Failure* 2013;1:31–9.
- Aslam S, Dan J, Topik A, et al. Decrease in driveline infections with change in driveline management protocol. *The VAD Journal* 2016;2.
- Hernandez RE, Singh SK, Hoang DT, et al. Present-Day Hospital readmissions after left ventricular assist device implantation: a large single-center study. *Tex Heart Inst J* 2015;42:419–29.
- Martin B-J, Luc JGY, Maruyama M, et al. Driveline site is not a predictor of infection after ventricular assist device implantation. *Asaio J* 2018;64:616–22.
- Raymond AL, Kfoury AG, Bishop CJ, et al. Obesity and left ventricular assist device driveline exit site infection. *Asaio J* 2010;56:57–60.
- Koval CE, Thuita L, Moazami N, et al. Evolution and impact of driveline infection in a large cohort of continuous-flow ventricular assist device recipients. *J Heart Lung Transplant* 2014;33:1164–72.
- John R, Aaronson KD, Pae WE, et al. Drive-line infections and sepsis in patients receiving the HVAD system as a left ventricular assist device. *J Heart Lung Transplant* 2014;33:1066–73.
- Trachtenberg BH, Cordero-Reyes A, Elias B, et al. A review of infections in patients with left ventricular assist devices: prevention, diagnosis and management. *Methodist Debakey Cardiovasc J* 2015;11:28–32.
- Imamura T, Murasawa T, Kawasaki H, et al. Correlation between driveline features and driveline infection in left ventricular assist device selection. *J Artif Organs* 2017;20:34–41.
- Haglund NA, Davis ME, Tricarico NM, et al. Readmissions after continuous flow left ventricular assist device implantation: differences observed between two contemporary device types. *Asaio J* 2015;61:410–6.
- Hannan MM, Xie R, Cowger J, et al. Epidemiology of infection in mechanical circulatory support: a global analysis from the ISHLT mechanically assisted circulatory support registry. *J Heart Lung Transplant* 2019;38:364–73.
- Gordon SM, Schmitt SK, Jacobs M, et al. Nosocomial bloodstream infections in patients with implantable left ventricular assist devices. *Ann Thorac Surg* 2001;72:725–30.
- Akhter SA, Badami A, Murray M, et al. Hospital readmissions after continuous-flow left ventricular assist device implantation: incidence, causes, and cost analysis. *Ann Thorac Surg* 2015;100:884–9.
- Bernhardt AM, Schlöglhofer T, Lauenroth V, et al. Prevention and early treatment of driveline infections in ventricular assist device patients - The DESTINE staging proposal and the first standard of care protocol. *J Crit Care* 2020;56:106–12.
- Girdler SJ, Glezos CD, Link TM, et al. The science of quality improvement. *JBJS Rev* 2016;4:e2.
- McCandless SP, Ledford ID, Mason NO, et al. Comparing velour versus silicone interfaces at the driveline exit site of HeartMate II devices: infection rates, histopathology, and ultrastructural aspects. *Cardiovasc Pathol* 2015;24:71–5.
- Dean D, Kallel F, Ewald GA, et al. Reduction in driveline infection rates: results from the HeartMate II multicenter Driveline silicone skin interface (SSI) registry. *J Heart Lung Transplant* 2015;34:781–9.
- Yoshitake S, Kinoshita O, Nawata K, et al. Novel driveline route for prevention from driveline infection: triple tunnel method. *J Cardiol* 2018;72:363–6.
- Cagliostro B, Levin AP, Fried J, et al. Continuous-Flow left ventricular assist devices and usefulness of a standardized strategy to reduce drive-line infections. *J Heart Lung Transplant* 2016;35:108–14.

- 33 Baronetto A, Centofanti P, Attisani M, *et al.* A simple device to secure ventricular assist device driveline and prevent exit-site infection. *Interact Cardiovasc Thorac Surg* 2014;18:415–7.
- 34 Kusne S, Mooney M, Danziger-Isakov L, *et al.* An ISHLT consensus document for prevention and management strategies for mechanical circulatory support infection. *J Heart Lung Transplant* 2017;36:1137–53.
- 35 Sousa-Uva M, Head SJ, Milojevic M, *et al.* 2017 EACTS guidelines on perioperative medication in adult cardiac surgery. *Eur J Cardiothorac Surg* 2018;53:5–33.
- 36 Chinn R, Dembitsky W, Eaton L, *et al.* Multicenter experience: prevention and management of left ventricular assist device infections. *Asaio J* 2005;51:461–70.
- 37 Barber J, Leslie G. A simple education tool for ventricular assist device patients and their caregivers. *J Cardiovasc Nurs* 2015;30:E1–10.
- 38 Nolan T, Resar R, Haraden C. *Improving the reliability of health care. IHI Innovation Series white paper.* Boston: Institute for Healthcare Improvement, 2004.
- 39 INTERMACS adverse event definitions: Adult and pediatric patients. INTERMACS website. UAB the University of Alabama at Birmingham website; STS InterMACS Database InterMACS Appendices - School of Medicine - Interagency Registry for Mechanically Assisted Circulatory Support | UAB.
- 40 Kirklin JK, Naftel DC, Stevenson LW, *et al.* INTERMACS database for durable devices for circulatory support: first annual report. *J Heart Lung Transplant* 2008;27:1065–72.
- 41 Benneyan JC, Lloyd RC, Plsek PE. Statistical process control as a tool for research and healthcare improvement. *Quality and Safety in Health Care* 2003;12:458–64.
- 42 Gupta M, Kaplan HC. Measurement for quality improvement: using data to drive change. *J Perinatol* 2020;40:962–71.
- 43 Schlöglhofer T, Michalovics P, Riebandt J, *et al.* Left ventricular assist device driveline infections in three contemporary devices. *Artif Organs* 2021;45:464–72.
- 44 Grimshaw JM, Shirran L, Thomas R, *et al.* Changing provider behavior: an overview of systematic reviews of interventions. *Med Care* 2001;39:112–45.
- 45 Wolfe HA, Taylor A, Subramanyam R. Statistics in quality improvement: measurement and statistical process control. *Paediatr Anaesth* 2021;31:539–47.
- 46 Wheeler TA, Davis JT, Brill R. The aggregate point rule for identifying shifts on P charts and U charts. *Pediatr Qual Saf* 2018;3:e103.