

# BMJ Open Quality Using clinical decision support tools to increase defibrillator deactivations in dying patients

Ramsey Kalil <sup>1</sup>, Daniel Y Choi,<sup>2</sup> Joshua D Geleris <sup>3</sup>, Jennifer I Lee,<sup>4</sup> Michael P Wagner<sup>5</sup>

**To cite:** Kalil R, Choi DY, Geleris JD, *et al.* Using clinical decision support tools to increase defibrillator deactivations in dying patients. *BMJ Open Quality* 2022;**11**:e001729. doi:10.1136/bmjopen-2021-001729

Received 16 November 2021  
Accepted 26 July 2022



© Author(s) (or their employer(s)) 2022. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

<sup>1</sup>Division of Cardiology, Texas Heart Institute, Houston, Texas, USA

<sup>2</sup>Division of Cardiology, New York University Medical Center, New York, USA

<sup>3</sup>Department of Medicine, NewYork-Presbyterian/Columbia University Medical Center, New York, USA

<sup>4</sup>Weill Department of Medicine, New York Presbyterian Hospital/Weill Cornell Medical Center, New York, New York, USA

<sup>5</sup>Department of Medicine, Memorial Sloan Kettering Cancer Center, New York, USA

## Correspondence to

Dr Ramsey Kalil;  
ramsey.kalil@gmail.com

## INTRODUCTION

Inadvertent automatic implantable cardioverter defibrillator (AICD) shocks at the end of life are distressing and should be avoided.<sup>1,2</sup> Up to 65% of AICDs are active and up to 33% of patients with AICDs receive shocks within 24 hours of dying.<sup>3</sup> Though guidelines encourage device discussions, one study demonstrated only 27% of clinicians brought up AICD deactivation.<sup>2,4</sup> Since the Health Information Technology for Economic and Clinical Health Act was passed in 2009, clinicians and IT professionals introduced evidence-based clinical decision support (CDS) models into patient care, including order sets and alerts.<sup>5</sup> Our group was the first to demonstrate that educational sessions paired with a novel CDS tool in the electronic medical record (EMR) significantly improved the rates of AICD deactivation discussions and of shock function deactivation.<sup>6</sup> However, a criticism was that the labour-intensive teaching sessions drove most of the improvement and EMR modifications are ineffective. As a result, we sought to investigate whether the CDS tool alone invoked more AICD deactivation discussions and AICD deactivations.

## METHODS

We conducted a retrospective chart review of hospitalised comfort care patients from April 2018 to April 2019 (n=46) at another academic medical centre within our hospital enterprise that uses the same EMR. The novel CDS tool was activated in October 2018 within the ‘comfort care’ order set, which directs providers

to focus on symptom-based treatment. Primary providers were prompted to address AICD deactivation planning by selecting one of three options in a drop-down menu—‘N/A patient does not have an AICD’, ‘patient or decision-maker declines AICD deactivation at this time. Do not deactivate AICD’, and ‘patient or decision-maker gives consent for AICD deactivation. Deactivate AICD’ (figure 1). No education or announcements of the CDS tool were made. The rates of AICD deactivation discussions (identified in a progress note) and AICD deactivations (presence of a deactivation note) were compared between the 6-month periods before (n=18) and after (n=28) CDS tool implementation. Fisher’s exact test was performed for comparison with a type I error cut-off of 0.05.

## RESULTS

Prior to implementing the CDS tool, 77% of patients underwent AICD deactivation discussion and 39% AICD deactivation (table 1). In 6-month postintervention, the rate of AICD deactivation discussions improved to 96%, nearing statistical significance (p=0.06). The rate of deactivated AICDs improved to 75% (p=0.02). Feedback from clinicians using this tool suggested that it was valuable and did not disturb standard workflow.

## DISCUSSION

In this study, we investigated whether a standalone CDS tool within the EMR could precipitate practice changes. Our data build on the

Per guidelines, discussion about AICD deactivation is required when a patient with an active AICD changes their goals of care to comfort care.	
This order documents that deactivation of the patient's AICD has or will be performed, however it is the ordering provider's responsibility to contact Cardiology/Electrophysiology directly to arrange for this deactivation.	
AICD Deactivation	
Order	Status
[-] AICD Deactivation - 1 item(s)	
<input checked="" type="checkbox"/> AICD Deactivation Planning	
Nursing General	N/A Patient does not have an AICD Patient or decision maker declines AICD deactivation at this time. Do not deactivate AICD. Patient or decision maker gives consent for AICD deactivation. Deactivate AICD.

**Figure 1** Screen capture of our group’s clinical decision support tool within the electronic medical record.

**Table 1** Improvement in ICD deactivations and ICD deactivation discussions after implementation of novel CDS tool in EMR\*

QI measure	Pre-CDS implementation	Post-CDS implementation	P value
ICD deactivation occurred	7	21	0.02
ICD deactivation did not occur	11	7	
ICD deactivation discussion occurred	14	27	0.06
ICD deactivation discussion did not occur	4	1	

\* Created by authors. \*\*Using Fisher's exact test.

CDS, clinical decision support; EMR, electronic medical record; ICD, implantable cardioverter defibrillator.

work of Sandhu and Matlock, who proposed a CDS tool to improve AICD deactivations, revealing that a properly implemented CDS tool adds value.<sup>7</sup> Similar tools have been successfully applied with associated improvement in quality metrics, including reduced length of hospital stay and fewer adverse drug events.<sup>8,9</sup>

Some argue against association of EMR-based CDS tools and better quality, raising concerns over their ability to improve healthcare delivery.<sup>10</sup> Inefficacy may reflect suboptimal design; missing or poor positioning of key elements can render a tool useless.<sup>11</sup> Alert fatigue may provoke failure to recognise areas to optimise healthcare performance.<sup>12</sup> We demonstrate improved outcomes through successful application of an intuitive CDS tool that prompts users to reconcile AICD deactivation planning at an appropriate juncture without disrupting workflow. Our tool was widely accepted and valuable without inundating providers with clicks, notifications or training modules, highlighting the importance of integration into existing systems. We conclude that continued provider education may be unnecessary to improve AICD deactivations in the presence of our CDS tool.

We note that 60% of patients who die at our institution receive comfort care orders, demonstrating a high yield for this CDS tool. Though another significant portion of deaths occur unexpectedly, these patients were not the targets of this intervention. In contrast, patients focused on comfort measures were appropriate candidates for AICD deactivation.

Limitations of our study include that it was conducted within a single hospital system, it lacked a control group, and there was a paucity of data on unintended AICD shock rates. Future directions involve analysing data about provider utilisation of the CDS tools, such as click patterns and time spent within an order set, to optimise design. Finally, expanding use of this CDS tool at other institutions and to patients who note a preference to not be resuscitated may encourage similar conversations to undergo upstream of end-of-life care.

**Twitter** Ramsey Kalil @RamseyKalilMD

**Contributors** The authors confirm contribution to the paper as follows: study conception and design: RK, JDG and MPW; data collection: RK and MPW; analysis and interpretation of results: RK, DYC, JDG, JIL, MPW; draft manuscript preparation: RK, DYC, JDG, JIL, MPW. All authors reviewed the results and approved the final version of the manuscript.

**Funding** The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

**Competing interests** None declared.

**Patient consent for publication** Not applicable.

**Ethics approval** The Office of Research Integrity conducted a review of the above-mentioned submission and determined that the activities described do not constitute human subjects research as the activities are not considered a systematic investigation designed to contribute to generalisable medical knowledge. Therefore, neither IRB approval nor a notice of exemptions are required to proceed.

**Provenance and peer review** Not commissioned; externally peer reviewed.

**Open access** This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

#### ORCID iDs

Ramsey Kalil <http://orcid.org/0000-0003-0668-838X>

Joshua D Geleris <http://orcid.org/0000-0003-0137-4111>

#### REFERENCES

- Fromme EK, Stewart TL, Jeppesen M, *et al*. Adverse experiences with implantable defibrillators in Oregon hospices. *Am J Hosp Palliat Care* 2011;28:304–9.
- Goldstein NE, Lampert R, Bradley E, *et al*. Management of implantable cardioverter defibrillators in end-of-life care. *Ann Intern Med* 2004;141:835–8.
- Stoevelaar R, Brinkman-Stoppelenburg A, Bhagwandien RE, *et al*. The incidence and impact of implantable cardioverter defibrillator shocks in the last phase of life: an integrated review. *Eur J Cardiovasc Nurs* 2018;17:477–85.
- Lampert R, Hayes DL, Annas GJ, *et al*. Hrs expert consensus statement on the management of cardiovascular implantable electronic devices (CIEDs) in patients nearing end of life or requesting withdrawal of therapy. *Heart Rhythm* 2010;7:1008–26.
- Health Information Technology for Economic and Clinical Health (HITECH) Act. *Title XIII of division A and title IV of division B of the American recovery and Reinvestment act of 2009 (arrA)*. 226. Pub L. No. 111-5, 123 Stat, 2009.
- Choi DY, Wagner MP, Yum B, *et al*. Improving implantable cardioverter defibrillator deactivation discussions in admitted patients made DNR and comfort care. *BMJ Open Qual* 2019;8:e000730.
- Sandhu A, Matlock D. Stopping superfluous shocks with system solutions. *JAMA Intern Med* 2020;180:1692–3.
- Macheel C, Reicks P, Sybrant C, *et al*. Clinical decision support intervention for rib fracture treatment. *J Am Coll Surg* 2020;231:249–56.
- Shahmoradi L, Safdari R, Ahmadi H, *et al*. Clinical decision support systems-based interventions to improve medication outcomes: a systematic literature review on features and effects. *Med J Islam Repub Iran* 2021;35:27–42.
- Romano MJ, Stafford RS. Electronic health records and clinical decision support systems. *Archives of Internal Medicine* 2011;171.
- Connor JP, Medow JE, Ehlenfeldt BD, *et al*. Electronic clinical decision support to facilitate a change in clinical practice: small details can make or break success. *Transfusion* 2020;60:1970–6.
- Shah SN, Amato MG, Garlo KG, *et al*. Renal medication-related clinical decision support (CdS) alerts and overrides in the inpatient setting following implementation of a commercial electronic health record: implications for designing more effective alerts. *J Am Med Inform Assoc* 2021;28:1081–7.