

BMJ Open Quality Resilience in nursing medication administration practice: a systematic review with narrative synthesis

Pollyanna LR Kellett ^{1,2}, Bryony Dean Franklin,^{3,4} Susie Pearce,^{2,5} Jonathan Benn¹

To cite: Kellett PLR, Franklin BD, Pearce S, *et al.* Resilience in nursing medication administration practice: a systematic review with narrative synthesis. *BMJ Open Quality* 2024;**13**:e002711. doi:10.1136/bmjopen-2023-002711

► Additional supplemental material is published online only. To view, please visit the journal online (<https://doi.org/10.1136/bmjopen-2023-002711>).

Received 9 December 2023
Accepted 13 September 2024



© Author(s) (or their employer(s)) 2024. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

¹School of Psychology, University of Leeds, Leeds, UK

²School of Nursing and Midwifery, University of Plymouth, Plymouth, UK

³UCL School of Pharmacy, London UK, London, UK

⁴Imperial College Healthcare NHS Trust, London, UK

⁵Torbay and South Devon NHS Foundation Trust, Torquay, UK

Correspondence to

Ms Pollyanna LR Kellett;
psplr@leeds.ac.uk

ABSTRACT

Resilience in nursing medication administration practice: a systematic review with narrative synthesis.

Objective Little is known about how nurses adapt medication administration practices to preserve safety. The capacity to adapt and respond before harm occurs has been labelled 'resilience'. Current evidence examining medication safety largely focuses on errors and what goes wrong. This review aimed to synthesise evidence for the application of resilience principles and practices in nursing medication administration.

Design The Preferred Reporting Items for Systematic Reviews and Meta-Analyses guided the review, which was registered with PROSPERO.

Data sources MEDLINE, EMBASE, PsychINFO and CINAHL databases were searched from 14 August 2020 to 1 January 2021 for English-language studies.

Methods A systematic review of empirical studies of any design relating to resilience and safety in nursing medication administration in the inpatient setting was conducted. Methodological quality was appraised using the Mixed Methods Appraisal Tool. Data were synthesised thematically.

Results Thirty-two studies with a range of methodologies of mostly good quality met the inclusion criteria. Eleven interventional studies included two that evaluated the effectiveness of education interventions and nine exploratory studies with outcomes showing the impact of an intervention designed or examined to build resilience. Twenty-one non-interventional studies showed how resilience principles are put into practice. Only three studies explicitly named the concept of resilience. Resilient medication administration strategies result from five triggers.

Conclusions Nurses' resilience practices were found to be responses to identified triggers that threaten safety and productivity. These were often short term, real-time proactive adaptations to preserve safety, compensating for and responding to complexities in the modern healthcare setting.

PROSPERO registration number CRD42018087928.

INTRODUCTION

Worldwide, registered nurses are accountable for preserving patient safety,¹ including the safe use of medicines.² Despite widespread efforts to improve safety through large-scale research, policy and regulation,³ it continues

WHAT IS ALREADY KNOWN ON THIS TOPIC

- ⇒ Resilience is a source of patient safety through adaptation to the constant variations and changing risks to maintain care quality and safety. Adaptations represent 'work-as-done' describing what actually happens in response to environmental complexity, as opposed to the 'work-as-planned' captured in guidance and standard operating procedures.
- ⇒ A focus on how safety is routinely maintained (ie, successful compensatory practice), as opposed to what goes wrong, is an important area of study for nursing medication administration practice.

WHAT THIS STUDY ADDS

- ⇒ Our analysis aligns the often short-term, real-time adaptive practices during nursing medication administration with the theoretical concept of resilience and identifies the triggers for 'work as done'.

HOW THIS STUDY MIGHT AFFECT RESEARCH, PRACTICE OR POLICY

- ⇒ This study strengthens conceptual and operational clarity for future research into nurses' capacity to adapt to dynamic risk to maintain safety in medication administration, paving the way for future interventions and strategies to promote resilience and patient safety.

to be a global concern.⁴ Medication administration is a key nursing function⁵ accounting for 54.9% of medication errors⁶ and taking between 11.8%⁷ and 29.1%⁸ of nurses' time.

The globally recognised framework for safe medication administration practice is the 'Five Rights' (5 R's), ensuring the right patient receives the right drug at the right dose, at the right time, via the right route; yet the 5 R's have received criticism for not acknowledging the human and systems factors inherent in medication administration.⁹

Traditionally safety is defined as absence of accidents and 'freedom from unacceptable risk', what is termed a 'safety-I' approach.¹⁰ More recently the concept of safety has been developed to include 'safety-II'. Here, the focus is not just risk and error prevention

but on preservation of safety under varying conditions by adapting to anticipated and emerging risks, termed ‘system resilience’.¹⁰ It is important to understand and design systems and practices that are more resilient to these dynamic risks¹¹; the nursing challenge being to promote, support and learn from responsive adaptations that maintain safety during medication administration. Everyday practice can be represented as ‘work as done’ reflecting the complex nature of clinical work, in contrast to ‘work as planned or imagined’ set out in policies and guidelines.¹²

Nurse’s adaptive or compensatory practices can be a key source of resilience when they preserve safety¹³; so understanding how nurses adapt to create safety and how resilience principles and practices are applied during nursing medication administration are important areas for enquiry.

The review aims to synthesise evidence for the application of resilience principles and practices in nursing medication administration by answering two specific research questions; how might resilience principles be applied? and what is the evidence for the effectiveness of strategies linked to resilience principles in improving safety in nurses’ medication administration? Interpretation of review findings is theoretically informed by two complementary frameworks. Holden *et al*’s¹⁴ Systems Engineering Initiative for Patient Safety (SEIPS) V.2.0 model, a human factor framework for understanding how care processes, individual and sociotechnical work shapes patient safety outcomes in resilient systems; and Aase *et al*’s¹⁵ framework

examining resilience emerging from individuals (internal resources) and organisations (external resources).

METHODS

Design

A systematic review was conducted following Centre for Reviews and Dissemination¹⁶ and PRISMA¹⁷ reporting guidelines and checklist. The Synthesis Without meta-analysis (SWiM) reporting items were used for quantitative interventional studies using alternative synthesis methods.¹⁸ The protocol was registered with PROSPERO.

Search methods

Initial scoping searches of Ovid Medline refined the search strategy. A faceted approach¹⁹ with three facets, ‘patient safety’ (including resilience synonyms), ‘nurse medication administration’ and ‘hospital inpatient’, was utilised. A context-specific definition of resilience was applied to the search, where resilience refers to adaptive or compensatory behaviour to maintain safe administration of medication in the face of variations and dynamic risks. For eligibility criteria, see [table 1](#).

Search strategy

Four databases were searched from 14 August 2020 to 1 January 2021 to find relevant studies published before 1 January 2021 in four databases: MEDLINE, EMBASE, PsychINFO and CINAHL. Boolean search terms were developed with assistance from a university subject librarian, the final strategy is shown in [box 1](#).

Table 1 Eligibility criteria

Context	Include	Exclude
Population	Registered nurses, registered nursing associates	Nursing students, any other healthcare professionals
Context	Administration of any medication, secondary care setting, any inpatient hospital setting, simulation of inpatient setting	Studies of drug incompatibilities, studies that examine incident reporting, any primary care setting, outpatients, day clinics, community drop-in centres, individual psychological resilience, studies using secondary analysis of existing data unless conducted by the original study team
Study focus	Patient safety, resilience (the ability to adjust functioning prior to, during, or following changes and disturbances in nursing medication administration practices to maintain high quality care)	Studies only reporting prevalence, measurement and characteristics of errors, studies only reporting reliability (the consistent, safe administration of medication), studies where nursing medication administration was not the main focus of the research or where data relating to nursing medication administration could not be extracted from the results
Country	Any country worldwide	
Publication type	English language, peer-reviewed studies of any study design meeting definition of resilience	Systematic reviews, non-empirical grey literature, conceptual articles, letters, non-English language, PhD and conference abstracts, PhD theses

Box 1 Example of search strategy (Ovid Medline)

```
(Patient safety or safety II or safety-II or safety 2 or (safety adj2 (culture or climate)) or (reliab$ or resilien$) or (human adj2 (factor$ or error$)) or (workaround$ or work around$ or work-around$ or circumvention or non-compliance or improvise$ or shortcut$ or double check$ or Safety Management/or Medication Errors/ or (error adj3 (detect$ or wisdom)).mp
AND
(((Nurs$ or Nursing) and ((medication$ or medicine$ or drug$ or dose or dosage or dosing) adj5 administration)).mp
AND
((in-patient$ or inpatient$ or acute or secondary care or hospital$ or Inpatient/ or Hospitals/)).mp
```

Study selection

Study identification was undertaken with two stages of inter-reviewer reliability, confirmation of full-text inclusion and quality assurance (see [figure 1](#)). A first-stage eligibility assessment of title and abstract by the primary author was confirmed for 5% of records with two reviewers confirming 2.5% each, with 97% agreement. A second stage title and abstract screen classified the records as empirical or non-empirical and confirmed the research focus with 95% confirmation of 5% of records with the reviewers. In both stages, unresolved records went through to full text review, an additional reviewer was not required. Further exclusions were made on subsequent examination of the full texts as some abstracts did not report sufficient detail, study type or focus. Final included studies were grouped according to whether or not they evaluated an intervention.

Quality appraisal

Concurrent quality appraisal used the Mixed Methods Appraisal Tool (MMAT),²⁰ appropriate for assessing a variety of methodologies and designs. No studies were excluded based on quality,²⁰ which was reported descriptively and confirmed by two reviewers for 30% of included studies.

Data extraction

Data were extracted and recorded using a bespoke tool developed through author piloting and review, displaying aspects of core details, study background and results. The detailed information is displayed in online supplemental files tables 1 and 2.

Synthesis of results

Meta-analysis of results across an extensive array of data was not possible due to heterogeneity in study outcomes and methodologies. Synthesis drew on the interpretive tradition, enabling maintenance of the rigour of the PRISMA and SWiM reporting guidelines and an iterative and dynamic approach to result synthesis. Data from result synthesis are provided (online supplemental table 2). A process of recurring theme identification grouped disparate data²¹ using an iterative three-stage approach

of coding, descriptive themes and analytic themes.²² This generated descriptive material to answer the first review question, and the analytic themes for the second. As part of the screening process, the included studies were differentiated by whether there was an interventional study design or not, but during result synthesis, this distinction became less discrete as the iterative themes evolved.

Patient and public involvement

As a PhD systematic review study, direct patient involvement in conduct of the research was not feasible. However, in initially proposing the topic, there was a keen awareness of the importance of the issues of medication safety and resilience in nurse medication administration practice for patients.²³ The systematic review forms the basis of a broader programme of work in this area by the authors, that is investigating how nurses adapt their administration practice to preserve patient safety, linked to two national research centres with medication safety research aims informed by patient advisory groups.

RESULTS

Identification of selected studies

The database search yielded 5416 records. Following screening, data were extracted from 32 studies. Identification of each stage of the search and selection process is summarised in the PRISMA diagram ([figure 1](#)).

Study characteristics

Included studies were published between 2005 and 2020 from five developed countries. The majority, 16 (50%) were conducted in the USA, with the remainder from Australia, 9 (28%), the Netherlands, 4 (13%), UK 2 (6%) and Norway, 1 (3%). Clinical specialities varied, with the majority, 22 studies (70%) in the adult setting, 8 (25%) in paediatrics, 1 (3%) was set in both adult and paediatric settings and 1 (3%) in a simulated clinical setting. Further study characteristics are outlined in online supplemental table 1. Although all studies cited core safety science literature related to resilience, only three studies explicitly referred to ‘resilience’ within the full text of their published articles.^{13 24 25}

Study methods and quality of evidence

Study methods varied (see online supplemental table 1). Applying the MMAT’s five categories of study design,²⁰ 22 (69%) studies had a qualitative design, 1 (3%) was a randomised controlled trial, 3 (9%) had quantitative non-randomised designs, 4 (12.5%) were quantitative descriptive and 2 (6%) were mixed method in design. The studies with qualitative designs often had more than one primary method of data collection, the majority using observational data, 7 (22%) or interview data, 6 (19%); and others using both observation and interviews, 4 (13%), focus groups, 3 (9%), and one study (3%) using observation, interview and focus groups; with another single study (3%) using focus groups and debriefs as primary methods to collect data. Quantitative non-randomised studies, 3




Figure 1 PRISMA flow chart. PRISMA, Preferred Reporting Items for Systematic Reviews and Meta-Analyses.

(9%) also used a variety of data collection techniques, including reports, 1 (3%), survey, 1 (3%) and observation with error reports, 1 (3%). The studies with a quantitative

descriptive design mostly used surveys, 3 (9%) with one study (3%) using observation to gather data. The single randomised control trial was an experimental simulated

study using observation; and the mixed method studies both combined qualitative with quantitative descriptive approaches, with one using observation with participant narratives and the other combining focus groups and surveys to gather data.

Methodological quality is rated from the lowest quality 0, through 25, 50, 75–100, the highest quality. The majority of studies were rated high quality (online supplemental table 1). The lowest scoring design (in the quantitative non-randomised category) scored 25. All the other categories of study design were rated high quality scoring 75–100. Certainty of the evidence for the findings from the 11 evaluative studies is described as per study authors and the methodological quality tool.²⁰ Methodological quality was rated consistently high in the qualitative evaluative studies. For the four quantitative studies, this was more varied, with one of lower quality,²⁶ two studies of very good quality^{27 28} and one of highest quality.²⁹

Findings

Thematic analysis enabled linkage between practices and strategies, and theoretical principles of resilience resulting in five conceptual themes. Two practices—workarounds and double checking spanned two themes. The 11 interventional studies included 4 quantitative and 7 qualitative studies; online supplemental table 2 outlines their interventions, effectiveness measures and results. Study quality for the quantitative studies varied (online supplemental table 1); their metrics for outcomes being reported as provided by study authors due to heterogeneity in design. Three of the four studies, one RCT²⁸ and two non-randomised designs^{26 29} evaluated the effectiveness of education interventions in respectively reducing interruptions and errors, improving scanning compliance and investigating the impact of work design on responses to missing medications. The fourth, a quantitative non-randomised study²⁷ assessed the impact of bar code medication administration on nurse's mental workload, their perceptions of safety and error likelihood. The remaining seven qualitative studies with interventions were of consistently high quality and were more exploratory in nature with outcomes showing the impact of an intervention designed to build resilience, or explore its use.

1. Resilience as a response to the introduction of technologies in the workplace

Workarounds were often cited as being detrimental to patient safety, while eight studies illustrated workarounds where nurses adjusted their behaviour when interacting with technology in the interest of preserving safety. These studies explored a range of outcomes, resulting from the introduction of medication-related technology (online supplemental table 2) such as identifying nurses' use of an electronic medication management system with some suggested targeted interventions³⁰; to documenting nurses' thinking processes³¹ problem-solving behaviours,^{24 32} perceptions of safety²⁷; frequency or causes of workarounds³³; or variability in practice³⁴ and

process problems.³⁵ Examples of resilience workarounds were nurses adapting technology-related protocols to ensure that a patient receives emergency medication for extremely low potassium³¹ and responding to wristband failure, scanner or bar code on medication failures by workarounds to adjust the system.³³ Other workarounds involved technology process issues, such as programming an intravenous pump for extra volume to be administered to make sure all the medication is given,^{32 34} nurses turning on the medication scanner in advance of use to account for slow booting or returning to paper-based records.²⁴ Here, nurses manually wrote down verbal medication orders, later following up with a call to physicians to issue an electronic version³⁵ or used paper-based records to 'mediate' the technology.³⁶ Contextual and human factors were found to play a role in perceptions of medication safety, linked by the 'fit' between the technology system and the nature of nurses' work²⁷ such as using the option in the barcode medication administration system to virtually extend a patient's hospitalisation to ensure analgesics can be administered to a patient in pain.³⁷

2. Resilience enhanced by education

Three nurse education programmes operationalising theoretical resilience principles were found (online supplemental table 2). The first study³⁸ described a medication safety programme aimed at increasing sensitivity to error risk and reduce medication administration errors, including case studies, simulation, meditation and a checklist. The second study introduced safety measures including lit lanyards, making the medication room a No Interruption Zone, and using the electronic medication record during handover to reduce interruptions.²⁶ The third study, a nursing leadership initiative, refined a scanning reporting system and educated nurses about their scanning habits to increase compliance with medication scanning.²⁹ Two of the education programmes reported only short-term (less than 3 months) reductions in medication errors.^{26 38} The first found an overall reduction in errors of 0.23% ($p < 0.003$) of total number of medications administered per day³⁸; the second found that errors reduced by 28 incidents over 3 months, compared with the same timeframe the previous year.²⁶ The third reported improvements in medication scanning compliance and other nursing safety practices, for example, identifying and correcting technical problems and making changes in clinical workflow to ensure patients receive timely medication prior to physiotherapy.²⁹

3. Resilience through critical thinking and problem-solving

The literature demonstrated resilience through nurse's critical thinking in response to a range of triggers. Decision-making was tested in a simulated setting²⁸ in response to two operational failures: missing medication from the medication cabinet and lack of insulin syringes. Here, nurses were found to choose a safer policy-compliant workaround rather than an unsafe one



(using a syringe not designed for insulin) only when they had access to a manager and when given the right work design (closeness to pharmacy to get the missing supplies). Under these conditions, nurses also contributed improvement ideas to enhance the future resilience of the system.²⁸ Constant professional vigilance was found to remain unchanged after barcode implementation,³¹ however the introduction of technology did result in new problem-solving behaviours. Here, a nurse used manual selection rather than the tethered scanner to identify a patient who was too far away to scan, and collaborated with a pharmacist to modify an ‘immediate’ prescription when the ordered dose of 20 mg was only available in two 10 mg doses, which they were unable to document as such on the system.²⁷ In exploring perceptions of safety, it was found that rather than their mental effort, nurses perceived it was how they reacted to external pressures such as feeling rushed, which impacted safety.²⁷

Problem-solving workarounds (violations and/or adaptations to policy) during medication administration were reported in two studies.^{39 40} When the reality of the environment was not matched by policy assumptions of the environment,³⁹ the intention to violate was influenced by the step of the administration process and triggered by whether it was a routine or emergency situation. Increased use of workarounds was linked to emotional exhaustion, but satisfaction with the administration system could exacerbate or reduce workarounds.⁴⁰

4. Responding to distractions and interruptions

Study authors in seven studies identified an array of theoretical principles of resilience in response to distraction and interruptions. These included ‘prioritisation’ to address more urgent need,^{25 41} ‘blocking’ to stop the interruption or ‘engagement’ with it,^{42 43} ‘prevention’ of any potential interruptions,^{42 44} ‘multitasking’⁴³ and ‘mediating’ by asking colleague to intervene.⁴³ Nurses have also been found to generate ‘personal safety systems’ such as using sticky notes or marking on paper medication cups to maintain safety when interrupted or facing compelling work demands.³⁶ In an attempt to avoid distractions on a busy unit, nurses found a quiet space to prepare medications before administration.⁴⁵ Those nurses who were more ‘task focused’ tended to minimise interruptions and distractions⁴⁴ by putting up a screen at the entrance of the clinical area. For predictable interruptions, found nurses used preventive strategies such as timing their medication rounds to avoid ward rounds or taking patients to the toilet prior to medication rounds.⁴² In unpredictable (usually emergency) situations, nurses most often used the strategy of ‘engagement’ with the interruption by prioritising it. This was usually to avoid compromising wider patient safety, for example, when a colleague asks for help with a baby having difficulty breathing.⁴²

5. Responding to uncertainty or novel situations

When responding to uncertainty or novel situations, three subthemes were identified.

Augmenting decisions

This was used to preserve safety where nurses gave greater attention to medications perceived as higher risk, for example, insulin and chemotherapy,^{13 46} and when handling time pressure by clustering medications prescribed for 8:00 or 10:00 by giving them together at 9:00 to facilitate timely administration.⁴⁷ Nurses used a smartphone light to check medication expiry dates in dimly lit areas, and an alarm reminder on their phone to ensure safety and timely medication administration.² Nurses were influenced by professional knowledge and patient data to ‘tinker’ and ‘tailor’ medication to individual patient need⁴⁸; and decisions to alter prescribed timings in response to the individual’s need for pain relief³⁷ or showing patients the medication chart to confirm that medications were appropriate⁴⁹ are resilient practices supporting safety.

Enhancing communication

Communication strategies to preserve safety included initiating and querying medication orders,⁵⁰ suggesting new medication along with the use of declarative language and closed questions,⁵¹ using open communication and verification, such as asking ‘is this what you want?’ to gain clarity,⁴⁵ and scanning the medication administration record during an informal verbal handover.⁵² There was some nuance between communication and double checking subthemes.

Double checking

As a resilient practice (performance of additional checks to those required by policy) is context specific and serves to maintain safety. Examples are nurses asking colleagues to double check²⁵ or even triple check medication not routinely used,⁴¹ inviting patients to verify medicines⁴⁴ or deciding to do another check of the patient and the drug⁵³ to alleviate uncertainty and promote safety.

6. Reported changes and suggestions for improvement

The included studies made various suggestions for change and future practice, signposting strategies supporting safety-II thinking themed as environment, resources and professional development. Within the worked environment future suggestions included roll out of a piloted safety programme,³⁸ using a medication safety care bundle to reduce interruptions,²⁶ ensuring technology reflects nurse’s tasks or context or usage,^{27 32 34 35} system redesign to support safety³⁹ and superusers to provide support.⁴⁰ Resource-focused recommendations were changes to design and usability of medication administration technology,^{24 27} increasing the availability of technology³⁰ and staff^{30 46}; stressing the importance of both understanding the complexity of medication administration,^{2 36 47} and the impact of policy on clinical judgement, to the organisation.⁵³ Suggestions for nurse professional

development centred on role modelling,³⁰ communication,⁵² including understanding of professional role and identities,⁵¹ enhancing insight recognising that safety extends beyond rules and procedure requiring critical thinking and judgement as part of professional expertise^{25 26 31 45 48 49}; and education initiatives.^{13 28 29 33 37 41–44 50}

DISCUSSION

This review aimed to answer two questions on the application of resilience practice in nursing medication administration—how might resilience principles be applied, and how effective are they in improving safety? Despite limited direct use of the concept of resilience in medication administration literature, this review has described and linked adaptive clinical practices (‘work-as done’) to resilience. Assessing effectiveness of resilience adaptive practices was only possible when this was an identified outcome measure, such as for educational interventions.

Evidence for the effectiveness of resilience strategies

Two intervention themes identifying external resilience (technology and education) and internal resilience (critical thinking) acknowledge the application of a resilience framework.¹⁵ Findings sensitive to interventions included identifying barriers to the use and acceptance of technology, measuring sensitivity to error risk and improving error rates, perceptions of safety and changes in problem-solving behaviours. Clear outcome and effectiveness measures in terms of safety were less clear, but they did generate understanding and findings relevant to safety in nurse medication administration. It is suggested that resilience is not tied to measures or estimates of probability of adverse events⁵⁴ and this was found to be the case in this examination of the effectiveness of resilience strategies. Only one study directly addressed resilience as a response to operational failure, demonstrating nurses’ capacity to adapt.²⁸ Two education strategies had short-term effects on error reduction^{26 38} but longer term follow-up would have further supported the significance of these studies. The impact of technology on nurses’ operational practice during medication administration warrants further exploration particularly regarding the necessity for workarounds, which demonstrate that some current configurations of technology/work system design are cumbersome and could be better optimised to support the human work of nurses.

‘Workarounds’ as resilience strategies

This review found examples of workarounds as either a response to working with technology or a response triggered by a situational context involving optimising decisions or critical thinking. Workarounds were interpreted by review authors as demonstrations of resilient solutions that promote safety, compensating for risks in the complex or unwieldy systems in which nurses administer medication. An alternative view sees workarounds as detrimental to patient safety, a source of risk and unstandardised variation. Insights from this analysis show

workarounds are innovative, compensatory behaviours triggered by context and have a role in preserving safety.

Interpretation of resilience in the work system

The findings of this review identify resilient practices, which serve to control the environment when responding to complexity and suggests nurses make choices in their adaptive practices depending on the context and triggers. For example, by blocking (ignoring) an interruption, nurses chose to prioritise administering medication. Yet most of the time nurses engaged with interruptions, stopping the primary task of medication administration.⁴³ This is considered to be an adaptive strategy that may support a broader system resilience by accepting a reduction in the quality of medication administration. Significantly, the application of aspects from the SEIPS V.2.0 model¹⁴ allowed description of the application of resilience practices (when either anticipated or unanticipated) as largely reactive, short-lasting intermittent adaptations; consistent with the work of Tucker and Edmondson⁵⁵ and similar to Hollnagel *et al*’s⁵⁶ ‘find and fix’ solutions to operational problems. An implication of the short time frame for these practices is that resilience is not stabilised and integrated in everyday work over time, and this misses the opportunity to achieve a planned, consistent, and high reliability environment. This has consequences for the safety of nursing medication administration. Findings from the studies in the review also highlight that adaptations can occur when available standardised protocols do not match the reality of clinical work. These variations in practice can be seen as vehicles for maintaining safety.^{13 39} Greater clarity and understanding of resilient medication administration practices have shown that they are often short-term, real-time adaptations to preserve safety, compensating for, and responding to, the complexities of working in the modern healthcare setting.

The resilient practices identified in this paper show how nurses respond to the need to manage. Their ability to adapt, anticipate and learn has clear implications for an alternative view of nurses’ safety strategies where the nurse is proactive in safeguarding the system, described as the nurse being the ‘Hero’, rather than the ‘Hazard’.⁵⁷ This aligns with safety-II thinking, with its focus on how safety and reliability are routinely maintained. This review has explored the nuance of how resilience in the medication administration context is defined, and the paper has identified a need for a stronger conceptual basis for intervention and evaluation in this area. Using suggestions from the resilience in healthcare programme,⁵⁸ an outcome from this review proposes the following definition of resilience in nursing medication administration for future research in this area:

‘Physical or mental compensatory or adaptive practices responding to detection of risks to patient safety before, during or after the process of administering medications to patients, where the nurse changes or



deviates from the planned model of work with the purpose of maintaining safety’.

Strengths and limitations

This review provides a comprehensive synthesis of the available evidence exploring resilient strategies in nursing medication administration. A major limitation was the lack of a standard definition of resilience as it applied to the context of nursing medication administration, with the possibility that with a broad concept like resilience, the search was not sensitive to all related work despite the application of a well-tested search strategy.

As encountered in this study, the broad heterogeneity in research focus and methods, differences in timeframes, scope of practice, technologies and policies across international data, pose challenges in terms of the practicality of a common conceptual framework for nursing resilience practice. However, it was possible to develop, through narrative synthesis, common themes in terms of broad resilience strategies and areas for improvement. Although the findings from the two large-scale studies set in the UK inpatient setting^{13 44} appear to confirm similarities in the application of resilient strategies, earlier research does not describe resilient behaviour per se, no doubt due to the relatively recent uptake of the concept in the patient safety and health services research literature. As such, findings of this review should be interpreted in light of the difficulties of post hoc application of the concept of resilience where it was not an explicit focus of the original study authors and lack of prior intention for identified practices, not always possible to infer, but a necessary part of describing the evidence.

Future research

Further alignment of resilient practices arising from medication administration with current thinking regarding the strengths of a safety-II approach and examining the tension between professional accountability and resilience are important to understand.

Future research might productively build on these review findings to better establish the concept of resilience and its application in administration practice. Exploring linked notions such as ‘work as done’ alongside policy (‘work as imagined’) is needed to support the design, evaluation and reporting of interventions that contribute to wider organisational safety.¹²

Once applied frameworks for resilience in this setting are in place, there is a need to establish the effectiveness of resilience strategies in medication administration on patient safety with robust interventional designs, clear outcome measures relating to safety improvement; and outcome measures that better represent resilience in the system in terms of nurses’ capacity to adapt. Further understanding of how resilience is achieved and how it can be sustained is key to developing resilient healthcare. This could

be supported by research to suggest ways to translate temporary resilience strategies into reliable error recovery ‘safety nets’ over time, potentially leading to improved medication administration safety.

CONCLUSION

This review set out to explore resilience in the context of nursing medication administration. Developing the understanding of how medication practices align with contemporary safety science theory has enabled description of the ways in which nurses exhibit resilient behaviours to maintain system safety in response to a number of contextual and environmental triggers that have been classified. These findings have implications for practice, policy and research and will strengthen conceptual and operational clarity by guiding the design, evaluation and reporting of interventions for supporting nurses’ capacity to adapt to dynamic risks in the complex systems in which they work.

Contributors PLRK, BDF, JB (guarantor): conceptualisation, methodology, validation, formal analysis, investigation, writing—original draft, writing—review and editing, visualisation, supervision, project administration. SP: formal analysis, investigation, writing—original draft, writing—review and editing, visualisation, supervision, project administration.

Funding PK is part-funded by University of Plymouth for PhD study, without external funding. BDF is supported by the National Institute for Health Research (NIHR) North West London Patient Safety Research Collaboration. SP has no external funding. JB is supported by the National Institute for Health Research (NIHR) Yorkshire and Humber Patient Safety Research Collaboration. The views expressed are those of the authors and not necessarily those of the NHS, the NIHR or the Department of Health and Social Care.

Competing interests The authors declare no competing interests.

Patient and public involvement Patients and/or the public were not involved in the design, or conduct, or reporting, or dissemination plans of this research.

Patient consent for publication Not applicable.

Ethics approval Not applicable.

Provenance and peer review Not commissioned; externally peer-reviewed.

Data availability statement Data sharing not applicable as no datasets generated and/or analysed for this study.

Supplemental material This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

ORCID iD

Pollyanna LR Kellett <http://orcid.org/0000-0001-7850-1356>

REFERENCES

- 1 Vaismoradi M, Tella S, A Logan P, et al. Nurses' Adherence to Patient Safety Principles: A Systematic Review. *Int J Environ Res Public Health* 2020;17:2028.
- 2 Martyn JA, Paliadelis P, Perry C. The safe administration of medication: Nursing behaviours beyond the five-rights. *Nurse Educ Pract* 2019;37:109–14.
- 3 Slawomirski L, Klazinga N. The economics of patient safety, from analysis to action oecd health working papers, no. 145. OECD Publishing, 2022. Available: <https://doi.org/10.1787/761f2da8-en>
- 4 World Health Organization. *Global patient safety action plan 2021–2030: towards eliminating avoidable harm in healthcare*. Geneva, Switzerland: World Health Organization, 2021. Available: <https://www.who.int/publications/i/item/9789240032705>
- 5 Shepherd M, Shepherd E. Medicines administration 1: understanding routes of administration. 2020; 116:42–4.
- 6 Elliott RA, Camacho E, Jankovic D, et al. Economic analysis of the prevalence and clinical and economic burden of medication error in England. *BMJ Qual Saf* 2021;30:96–105.
- 7 Michel O, Garcia Manjon A-J, Pasquier J, et al. How do nurses spend their time? A time and motion analysis of nursing activities in an internal medicine unit. *J Adv Nurs* 2021;77:4459–70.
- 8 Keohane CA, Bane AD, Featherstone E, et al. Quantifying nursing workflow in medication administration. *J Nurs Adm* 2008;38:19–26.
- 9 Institute for Safe Medication Practices (ISMP). The five rights: a destination without a map acute care ISMP medication safety alert! Institute for Safe Medication Practices (ISMP); 2007. Available: <https://www.ismp.org/resources/five-rights-destination-without-map>
- 10 Hollnagel ET. *Safety-II in practice: developing the resilience potentials*. Abingdon, Oxon: Routledge, 2017.
- 11 World Health Organization. Medication without harm - global patient safety challenge on medication safety. Geneva, Switzerland World Health Organization; 2017.
- 12 Ifaifel M, Lim RH, Ryan K, et al. *Resilient health care: a systematic review of conceptualisations, study methods and factors that develop resilience*. BMC Health Services Research, 2020.
- 13 Vos J, Franklin BD, Chumbley G, et al. Nurses as a source of system-level resilience: Secondary analysis of qualitative data from a study of intravenous infusion safety in English hospitals. *Int J Nurs Stud* 2020;102:103468.
- 14 Holden RJ, Carayon P, Gurses AP, et al. SEIPS 2.0: a human factors framework for studying and improving the work of healthcare professionals and patients. *Ergonomics* 2013;56:1669–86.
- 15 Aase K, Guise V, Billett S, et al. Resilience in Healthcare (RiH): a longitudinal research programme protocol. *BMJ Open* 2020;10:e038779.
- 16 Centre for Reviews and Dissemination. Systematic reviews: CRD's guidance for undertaking reviews in healthcare. 2009. Available: https://www.york.ac.uk/media/crd/Systematic_Reviews.pdf
- 17 Page MJ, McKenzie JE, Bossuyt PM, et al. The PRISMA 2020 statement: an updated guideline for reporting systematic reviews. *Br Med J* 2021;372:n71.
- 18 Campbell M, McKenzie JE, Sowden A, et al. Synthesis without meta-analysis (SWiM) in systematic reviews: reporting guideline. *BMJ* 2020;368:16890.
- 19 Fagan JC. Usability Studies of Faceted Browsing: A Literature Review. *ITAL* 2013;29:58–66.
- 20 Hong QN, Fàbregues S, Bartlett G, et al. The Mixed Methods Appraisal Tool (MMAT) version 2018 Education for Information. 2018;34:285–91.
- 21 Lucas PJ, Baird J, Arai L, et al. Worked examples of alternative methods for the synthesis of qualitative and quantitative research in systematic reviews. *BMC Med Res Methodol* 2007;7:4.
- 22 Thomas J, Harden A. Methods for the thematic synthesis of qualitative research in systematic reviews. *BMC Med Res Methodol* 2008;8:45.
- 23 Giles SJ, Lewis PJ, Phipps DL, et al. Capturing Patients' Perspectives on Medication Safety: The Development of a Patient-Centered Medication Safety Framework. *J Patient Saf* 2020;16:e324–39.
- 24 Holden RJ, Rivera-Rodriguez AJ, Faye H, et al. Automation and adaptation: Nurses' problem-solving behavior following the implementation of bar coded medication administration technology. *Cogn Technol Work* 2013;15:283–96.
- 25 Alteren J, Hermstad M, White J, et al. Conflicting priorities: Observation of medicine administration. *J Clin Nurs* 2018;27:3613–21.
- 26 Freeman R, McKee S, Lee-Lehner B, et al. Reducing interruptions to improve medication safety. *J Nurs Care Qual* 2013;28:176–85.
- 27 Holden RJ, Brown RL, Scanlon MC, et al. Micro- and macroergonomic changes in mental workload and medication safety following the implementation of new health IT. *Int J Ind Ergon* 2015;49:131–43.
- 28 Tucker AL. The impact of workaround difficulty on frontline employees' response to operational failures: A laboratory experiment on medication administration. *Manage Sci* 2016;62:1124–44.
- 29 Van Ornum M. Improving Bar Code Medication Administration Compliance in a Community Hospital Through a Nursing Leadership Initiative. *J Nurs Care Qual* 2018;33:341–7.
- 30 Debono D, Taylor N, Lipworth W, et al. Applying the Theoretical Domains Framework to identify barriers and targeted interventions to enhance nurses' use of electronic medication management systems in two Australian hospitals. *Implement Sci* 2017;12.
- 31 Eisenhauer LA, Hurley AC, Dolan N. Nurses' reported thinking during medication administration. *J Nurs Scholarsh* 2007;39:82–7.
- 32 McAlearney AS, Vrontos JJ, Schneider PJ, et al. Strategic Work-Arounds to Accommodate New Technology: The Case of Smart Pumps in Hospital Care. *J Patient Saf* 2007;3:75–81.
- 33 Rack LL, Dudjak LA, Wolf GA. Study of nurse workarounds in a hospital using bar code medication administration system. *J Nurs Care Qual* 2012;27:232–9.
- 34 Nemeth CP, Brown J, Crandall B, et al. The mixed blessings of smart infusion devices and health care IT. *Mil Med* 2014;179:4–10.
- 35 Niazkhani Z, Pirnejad H, van der Sijs H, et al. Evaluating the medication process in the context of CPOE use: the significance of working around the system. *Int J Med Inform* 2011;80:490–506.
- 36 Boonen M, Rankin J, Vosman F. Nurses' knowledge and deliberations crucial to Barcoded Medication Administration technology in a Dutch hospital: Discovering nurses' agency inside ruling. *Health (Lond)* 2020;24:279–98.
- 37 Boonen M, Vosman FJH, Niemeijer A. How bar coded medication administration technology affects the nurse-patient relationships: An ethnographic study. *Int J Technol Assess Health Care* 2018;34:507–13.
- 38 Durham ML, Suhayda R, Normand P, et al. Reducing Medication Administration Errors in Acute and Critical Care: Multifaceted Pilot Program Targeting RN Awareness and Behaviors. *J Nurs Adm* 2016;46:75–81.
- 39 Alper SJ, Holden RJ, Scanlon MC. Self-reported violations during medication administration in two paediatric hospitals. *BMJ Qual Saf* 2012;21:408–15.
- 40 Halbesleben JR, Rathert C, Williams ES. Emotional exhaustion and medication administration work-arounds: the moderating role of nurse satisfaction with medication administration. *Health Care Manage Rev* 2013;38:95–104.
- 41 Colligan L, Bass EJ. Interruption handling strategies during paediatric medication administration. *BMJ Qual Saf* 2012;21:912–7.
- 42 Johnson M, Weidemann G, Adams R, et al. Predictability of Interruptions During Medication Administration With Related Behavioral Management Strategies. *J Nurs Care Qual* 2018;33:E1–9.
- 43 Sitterding MC, Ebright P, Broome M, et al. Situation Awareness and Interruption Handling During Medication Administration. *West J Nurs Res* 2014;36:891–916.
- 44 McLeod M, Barber N, Franklin BD. Facilitators and Barriers to Safe Medication Administration to Hospital Inpatients: A Mixed Methods Study of Nurses. *Med Admin Processes Syst (MAPS Study) PLOS ONE* 2015;10:e0128958.
- 45 Dickson GL, Flynn L. Nurses' clinical reasoning: processes and practices of medication safety. *Qual Health Res* 2012;22:3–16.
- 46 Davis L, Keogh S, Watson K. Dishing the drugs: a qualitative study to explore paediatric nurses' attitudes and practice related to medication administration. *Collegian* 2005;12:15–20.
- 47 Jennings BM, Sandelowski M, Mark B. The nurse's medication day. *Qual Health Res* 2011;21:1441–51.
- 48 Boonen M, Vosman FJH, Niemeijer AR. Tinker, tailor, deliberate. An ethnographic inquiry into the institutionalized practice of bar-coded medication administration technology by nurses. *Appl Nurs Res* 2017;33:30–5.
- 49 Martyn JA, Paliadelis P. Safe medication administration: Perspectives from an appreciative inquiry of the practice of registered nurses in regional Australia. *Nurse Educ Pract* 2019;34:111–6.
- 50 Aydon L, Hauck Y, Zimmer M, et al. Factors influencing a nurse's decision to question medication administration in a neonatal clinical care unit. *J Clin Nurs* 2016;25:2468–77.
- 51 Borrott N, Kinney S, Newall F, et al. Medication communication between nurses and doctors for paediatric acute care: An ethnographic study. *J Clin Nurs* 2017;26:1978–92.
- 52 Braaf S, Rixon S, Williams A, et al. Medication communication during handover interactions in specialty practice settings. *J Clin Nurs* 2015;24:2859–70.



- 53 Davis L, Ware RS, McCann D, *et al.* Factors influencing paediatric nurses' responses to medication administration. *BMJ Qual Saf* 2010;19:e4.
- 54 Nemeth C, Wears R, Woods D. *Minding the gaps: creating resilience in health care*. USA: Rockville (MD): Agency for Healthcare Research and Quality (US), 2008.
- 55 Tucker AL, Edmondson AC. Why Hospitals Don't Learn from Failures: Organizational and Psychological Dynamics That Inhibit System Change. *Calif Manage Rev* 2003;45:55–72.
- 56 Hollnagel E, Wears RL, Bratihwaite J. From safety-I to safety-II: a white paper: the resilient health care net. University of Southern Denmark, University of Florida, USA, and Macquarie University, Australia; 2015. Available: <https://psnet.ahrq.gov/issue/safety-i-safety-ii-white-paper>
- 57 Reason J. *The human contribution – unsafe acts, accidents and heroic recoveries*. 1st edn. Aldershot, England: Ashgate, 2008.
- 58 Wiig S, Aase K, Billett S, *et al.* Defining the boundaries and operational concepts of resilience in the resilience in healthcare research program. *BMC Health Serv Res* 2020;20:330.