Abstract 1050 Figure 1 Percentage of type and screen drawn for laparoscopic hysterectomy procedure p chart, sigma

(21%), 5 patients (1%) received transfusion (Table 1). Figure-1 shows a process control chart plotting the T and S samples over time with interventions at 5–2015 and 4–2016 which show a significant decrease in mean T and S samples.

Abstract 1050 Table 1

	Laparoscopic Hysterectomy Baseline (1/2014 - 4/2015)	Laparoscopic Hysterectomy Intervention 1 (5/2015 - 3/2016)	Laparoscopic Hysterectomy Intervention 2 (4/2016 - 2/2017)
Total	615	490	552
Type and Screen	490 (78%)	300 (61%)	144 (26%)
Transfusion	21 (3%)	18 (3.6%)	5 (<1%)

Conclusions Unnecessary preparation of blood products for operations with historically low rates of transfusion represents wasted phlebotomy, labour, and expense. Using laparoscopic hysterectomy as an example, we decreased unnecessary T and S using data to guide pre-operative testing.

1057

REDUCING UNWANTED AND UNWARRANTED ED AND HOSPITAL UTILISATION FOR FRAIL ELDERLY IN RURAL SKILLED NURSING FACILITIES: A HYBRID IMPROVEMENT-IMPLEMENTATION APPROACH

¹Brant Oliver, ²Daniel Stadler, ²Ellen Flaherty, ¹Stephen Bartels. ¹Dartmouth Collaboratory for Health Care Delivery Science, US; ²Dartmouth Hitchcock, US

10.1136/bmjopen-2017-01129

Background Frail elders in residential skilled nursing facilities (SNFs) have high rates of emergency department (ED) utilisation and hospitalizations. We sought to implement and iteratively specify an intervention to improve utilisation and cost outcomes for frail elders in rural SNFs.

Objectives (1) To reduce unwanted and avoidable ED utilisation and hospitalizations; (2) to reduce related costs.

Methods Adopting evidence based practices, we iteratively developed an implementation approach including the following key elements: (1) advanced directives; (2) a dedicated closed-

call team of providers following SNF residents; (3) a biweekly case review of all ED referrals and hospitalizations; and (4) a standardised triage communication process. We conducted three PDSA cycles over a 6 month period and assessed clinical and cost outcomes using inferential statistics and statistical process control (SPC) methods.

Results Three rural SNFs participated in the intervention from January-June 2016. Three PDSA cycles were conducted. Monthly hospital-based care utilisation for long term care (LTC) residents reduced from 10 to 3.3 episodes ($p < 0.05$). ED transfers reduced by 59% ($p < 0.05$), and hospitalizations reduced by 62% ($p < 0.05$), without associated changes in

Timeline and PDSA Cycles

Abstract 1057 Figure 1

Abstract 1057 Figure 2

overall mortality or quality performance indicators. SPC analysis suggests that PDSA Cycle 2 was temporally related to the onset of improvement. Estimated cost savings calculated based on utilisation reductions is approximately \$7 million annually. **Conclusions** This initial pilot was feasible, achieved significant outcomes, and is an example of the use of improvement methods to iteratively develop and optimise an implementation approach. This approach has potential to significantly impact outcomes, utilisation and cost and is worthy of continued study.

Abstract 1057 Figure 3

1058 REDUCING EMERGENCY DEPARTMENT UTILISATION IN INFANTS: A QUALITY IMPROVEMENT STUDY

¹Jenny Chan, ²Mariam Maloyan, ²Hannah Durant, ²Ellen Reisinger, ²Pamela Schubert, ²Grace Chi, ²Joanne Cox, ²Kathleen Conroy. ¹Boston Children's Hospital Primary Care at Longwood, US; ²Boston Children's Hospital Primary Care at Martha Eliot, US

10.1136/bmjopen-2017-013130

Background Utilising the Emergency Department (ED) rather than the primary care medical home (PCMH) for non-emergent care results in decreased continuity and increased medical expense. Young, low-income children have the highest low-acuity ED visit rates. Few PCMH-based interventions have been shown to decrease unnecessary ED use.

Objectives Reduce rates of low-acuity ED utilisation in children under 12 months.

Methods This was a quality improvement study set in an academic primary care clinic serving 15,000 predominantly low-income families. Interventions focused on expanding urgent care (UC) and nursing-line access, improving parents' awareness of these ED-alternatives, and enhancing caregiver knowledge of infant care (Figure 1). Our primary outcome was the 12 month rolling rate of low-acuity ED visits; high-acuity ED visits acted as a balancing measure.

Abstract 1058 Figure 1 Driver diagram

Abstract 1058 Figure 2 ED utilisation by patients <1 year, 12 month rolling rate (low acuity visits)