Abstract 886 Figure 1 Statistical process control chart (p-chart) 3 sigma

increased to 87.9% (181/206) with 82.5% (n=170) independent, sustaining the improvement at >90% since October 2016.

Conclusions Ensuring families' independence with CL care in the home is a safety priority which can be achieved through a

teach-back program embedded into routine care. Determining impact on families' CL care-associated distress and ambulatory CLABSI rates is in process.

899 STOP MISSING ELEVATED BLOOD PRESSURES (BP): USING QI METHODOLOGY TO IMPROVE BP MANAGEMENT IN AN OUTPATIENT PAEDIATRIC NEPHROLOGY CLINIC

¹Hailey Woollen, ²Maureen Walsh Korick, ³Cheryl Courtlandt, ³Karen Emmerton, ³Mehgan Beverley, ³Susan Massengill. ¹Cincinnati Children's, US; ²Queens University of Charlotte, US; ³Levine Children's Hospital @ CMC, US

10.1136/bmjopen-2017-IHL8

Background Despite previous studies demonstrating accelerated deterioration in kidney function due to hypertension in paediatric patients with chronic kidney disease, there are significant discrepancies in the identification and management of elevated blood pressure in this high-risk population. Recognition of hypertension during childhood is hampered by the need for gender, age and height adjustment in calculations. We

STOP missing elevated blood pressures (BP): Using QI methodology to improve BP management in an outpatient nephrology clinic

Abstract 899 Figure 1 Percentage of clinic visits with STOP forms (p chart)

STOP missing elevated blood pressures (BP): Using QI methodology to improve BP management in an outpatient nephrology clinic

Abstract 899 Figure 2 Elevated BP percentile with intervention (p chart)

developed the Surveillance Target Outpatient blood Pressure (STOP) decision support tool to prompt team members to communicate BP percentile to identify hypertensive patients.

Objectives Improve recognition and intervention for elevated BP in ages 2–17 in a paediatric nephrology clinic. 1.) 90% of visits utilise STOP form 2.) 90% of elevated BP percentiles have documented intervention

Methods Model for Improvement framework was employed for this paediatric resident initiated QI project including rapid cycle PDSA testing beginning January through August 2017. Multidisciplinary teammates included receptionists, CMAs, nurses, and physicians. Daily data collection included quantitative measures analysed using run charts and control charts.

Results STOP form usage increased from an average of 69% to 92% (n=1498) using rapid cycle improvement by calculating and communicating BP percentile. Significant improvement was achieved in elevated BP management from an average of 74% to 100% of patients (n=483) with decreased variability occurring over the span of the project.

Conclusions We successfully used QI methods and tools to develop and implement a team-focused decision support tool to standardise and improve identification and management of elevated BP percentiles. The STOP form can be adopted and utilised in numerous clinical settings to identify those at risk for hypertension

902 REDUCED READMISSION RATES THROUGH MULTIDISCIPLINARY, WHOLE PERSON CARE

Mary Zozaya-Monohon, Andrea Corona. *Providence Health and Services, US*

10.1136/bmjopen-2017-IHL9

Background Readmissions are a surrogate marker for a patient's ability to manage their health. Ambulatory care is poised to transform services by partnering with patients to enhance their success with disease management to reduce hospitalizations.

Objectives Empower patients to optimise whole person, self-care with the support of a multidisciplinary team.

Methods A target population was selected after analysis of hospitalised and readmitted patients to confirm findings reported in the literature. A multidisciplinary team developed goals and objectives for whole person care to provide the framework for interventions. A process for identification, enrollment and discharge from the team was created. During

Abstract 902 Figure 1 Readmission rate run chart. The median readmission rate at baseline was 28% (20th, 80th percentiles 11,31) and decreased to 10% (20th, 80th percentiles 6,13).

team meetings patient needs are prioritised and integrated services are coordinated. The electronic medical record supports communication to the primary care provider and other members of the healthcare team.

Results The median readmission rate at baseline was 28% and decreased to 10% post implementation. This reduction in readmissions was obtained by implementing the intervention in 20% of all admitted patients. Interventions were assessed at baseline in all discharged patients compared to post-implementation target population. Completion of Physician's Orders for Life-Sustaining Treatment (POLST) rose from 9% to 41%, Personal Health Questionnaire-9 and Generalised Anxiety Disorder screening significantly increased to 94%. Evaluation by behavioural health and a pharmacist improved to 100%.

Conclusions Readmission rates can be decreased through coordinated, whole person interventions on a targeted patient population in the ambulatory setting. Implementing multidisciplinary proactive planned care improves a patient's ability to care for themselves as demonstrated by a reduced readmission rate.

Abstract 902 Figure 2 Percent of discharged patients receiving intervention

Abstract 902 Table 1 Interventions completed at baseline for all discharged patients compared to post-implementation in the target population

	All discharged patients at baseline (n=169)	Target-Population [^] post-intervention (n=94)
Completion of POLST	15 (9%)	38 (41%)
PHQ-9 Screening	14 (8%)	88 (94%)
GAD-7 Screening	0 (0%)	88 (94%)
Behavioural Health	15 (9%)	94 (100%)
Evaluation		
Pharmacist	7 (4%)	94 (100%)
Evaluation		

[^] Readmission within 90 days or a hospitalisation with a comorbid diagnosis of chronic obstructive pulmonary disease, diabetes, or heart failure.