

Importance of safety climate, teamwork climate and demographics: understanding nurses, allied health professionals and clerical staff perceptions of patient safety

Shahram Zaheer,¹ Liane R Ginsburg,¹ Hannah J Wong,¹ Kelly Thomson,² Lorna Bain^{3,4}

To cite: Zaheer S, Ginsburg LR, Wong HJ, et al. Importance of safety climate, teamwork climate and demographics: understanding nurses, allied health professionals and clerical staff perceptions of patient safety. *BMJ Open Quality* 2018;7:e000433. doi:10.1136/bmjoq-2018-000433

Received 16 May 2018

Revised 2 October 2018

Accepted 9 October 2018



Check for updates

© Author(s) (or their employer(s)) 2018. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

¹School of Health Policy and Management, York University, Toronto, Ontario, Canada

²School of Administrative Studies, York University, Toronto, Ontario, Canada

³Interprofessional Collaboration and Education, Southlake Regional Health Centre, Newmarket, Ontario, Canada

⁴University of Toronto, Toronto, Ontario, Canada

Correspondence to
Dr. Shahram Zaheer;
szaheer@yorku.ca

ABSTRACT

Background There is growing evidence regarding the importance of contextual factors for patient/staff outcomes and the likelihood of successfully implementing safety improvement interventions such as checklists; however, certain literature gaps still remain—for example, lack of research examining the interactive effects of safety constructs on outcomes. This study has addressed some of these gaps, together with adding to our understanding of how context influences safety.

Purpose The impact of staff perceptions of safety climate (ie, senior and supervisory leadership support for safety) and teamwork climate on a self-reported safety outcome (ie, overall perceptions of patient safety (PS)) were examined at a hospital in Southern Ontario.

Methods Cross-sectional survey data were collected from nurses, allied health professionals and unit clerks working on intensive care, general medicine, mental health or emergency department.

Results Hierarchical regression analyses showed that perceptions of senior leadership ($p<0.001$) and teamwork ($p<0.001$) were significantly associated with overall perceptions of PS. A non-significant association was found between perceptions of supervisory leadership and the outcome variable. However, when staff perceived poorer senior leadership support for safety, the positive effect of supervisory leadership on overall perceptions of PS became significantly stronger ($p<0.05$).

Practice implications Our results suggest that leadership support at one level (ie, supervisory) can substitute for the absence of leadership support for safety at another level (ie, senior level). While healthcare organisations should recruit into leadership roles and retain individuals who prioritise safety and possess adequate relational competencies, the field would now benefit from evidence regarding how to build leadership support for PS. Also, it is important to provide on-site workshops on topics (eg, conflict management) that can strengthen working relationships across professional and unit boundaries.

BACKGROUND

Empirical evidence suggests that an adverse event occurs in up to 10% of hospitalisations and that half of these events are preventable.¹

Healthcare errors are costly; for example, surgical site infections in Canadian hospitals result in \$24.4 million in extra healthcare costs annually.² Application of standardised clinical interventions such as hand hygiene guidelines³ and surgical checklists⁴ has been shown to reduce preventable medication and diagnostic and surgical errors. However, there is also a growing body of evidence that contextual factors such as safety climate influence the likelihood of successfully implementing safety improvement interventions—for example, checklists⁵ and initiatives to reduce central line infections.⁶

The objective of this paper is to examine the relationships (ie, direct and moderated) between staff (ie, nurses, allied health professionals (AHP) and unit clerks) perceptions of safety climate, teamwork climate and a self-reported safety measure. Safety climate was considered as senior and supervisory leadership support for safety.

Theory and literature review

According to high reliability theory (HRT), leaders shape safety climate perceptions of employees by prioritising or subordinating safety over other organisational/departmental goals such as efficiency. This in turn enables or discourages employee participation in safety enacting behaviours (such as teamwork and speaking up) that can improve safety outcomes and reducing failures at the frontlines. These enabling and enacting processes interact with each other to further strengthen/entrench safety climate and safety improvement gains.^{7,8}

Healthcare researchers have primarily relied on surveys to capture climate perceptions of safety. However, some of these surveys are very broad and contain antecedent or

outcome dimensions, in addition to *components* of safety climate.^{9 10} The construct of safety climate was originally defined as employee's perceptions of management commitment to safety¹¹ and the research community has been encouraged to adhere to this original definition of safety climate in order to maximise the practical utility of the construct.^{10 12} Indeed, perceptions of climate are domain specific, for example, safety climate, teamwork climate, innovation climate, and so on—rather than global or generic in nature.⁹ This domain-specific definition of climate also implies the simultaneous existence of a number of interacting climates in any given organisation. The current study adheres to this more focused definition and defines safety climate as employees' perception of leadership support for safety at two levels—senior and supervisory. Employees differentiate between the priorities of senior management and unit supervisors, resulting in the emergence of perceptions of two concurrent levels of safety climate.^{13 14} Adopting a multilevel safety climate perspective is especially important in loosely coupled organisations such as hospitals where unit supervisors can often exercise discretion in implementing policies created by senior management. Indeed, one key aspect of evaluating safety climate lies in examining consistency between organisational-level safety policies and implementation practices in subunits that are subject to supervisory discretion.¹⁴

In addition, much of the empirical research in healthcare settings has primarily focused on safety-related behaviours of leaders such as participative leadership style,¹² empowering leadership style,¹⁵ executive safety walk-rounds,^{16 17} adopt-a-work unit¹⁸ and front-line safety forums¹⁹ that can improve employees' perceptions of safety climate. However, empirical support for the predictive power of staff perceptions of leadership on safety outcomes remains limited. Empirical studies that have examined such relationships suggest that presence of supportive leaders significantly improves staff outcomes (eg, decreased nurse burnout)²⁰ and patient outcomes (eg, decreased falls, nosocomial infections, medication errors and adverse events).^{21 22} Only a handful of empirical studies have attempted to simultaneously evaluate the interactive impact of staff perceptions of senior and supervisory leadership on safety outcomes.²³ There is a need for further empirical work to more deeply examine how different levels of leadership may influence safety.

Teamwork climate also continues to be seen as an important predictor of safety outcomes. In the past, highly specialised professionals operating in silos were often sufficient to provide appropriate treatment to patients. However, changing disease patterns and growing complexity of care delivery now require teamwork behaviours including open communication²⁴ and semisynchronous cooperation²⁵ across providers and provider groups in order to reduce preventable errors and improve safety outcomes.²⁶ Indeed, emerging empirical evidence suggests that positive staff perceptions of teamwork climate are associated with reduced incidence of

in-hospital adverse events,²² lower rates of staff burnout,²⁷ reduced odds of poor surgical outcomes²⁸ and reduced patient mortality.²⁹ Lastly, empirical research in clinical settings also suggests that employees' perceptions of leadership first directly influences quality of teamwork and then through teamwork influences safety outcomes—for example, adverse events, burnout.^{20 27}

Justification for the current study

The research community has made important inroads in understanding the role of contextual factors in improving staff and patient safety (PS). However, the current study addresses several gaps that persist in the literature: (a) the need for more theory-based empirical research on safety-related contextual factors and a more focused definition of safety climate¹⁰; (b) a dearth of studies that examine the unique variance accounted for by multilevel perceptions of safety climate on a safety outcome²³; (c) the need for empirical research that examines the main as well as interactive effects of safety climate on outcomes³⁰; and (d) empirical research on safety and quality has primarily focused on understanding the perceptions of nurses while largely ignoring the perspectives of other healthcare professionals.

The aim of the current HRT informed study is to examine the influence (ie, main and interactive effects) of nurses, AHPs and unit clerks' perceptions of safety climate and teamwork climate on a self-reported safety outcome. More specifically, it is hypothesised that:

Hypothesis 1: Perceptions of senior leadership support for safety and supervisory leadership support for safety will both be positively associated with overall perceptions of PS. Furthermore, these predictor variables will interact and significantly influence the outcome variable.

Hypothesis 2: Perceptions of teamwork climate will be positively associated with overall perceptions of PS. Furthermore, teamwork climate and multilevel safety climate (ie, perceptions of leadership support for safety at the senior and supervisory levels) will have a significant interaction effect on outcome variable.

METHODS

Sampling and data collection procedures

The study was conducted at a large community hospital located in Southern Ontario (Canada). Survey data were obtained from front-line nurses (eg, registered nurses, registered practical nurses), AHPs (eg, respiratory therapists, physiotherapists, pharmacists) and clerical staff who worked on one of the four participating clinical units: intensive care unit (ICU), general medicine, adult inpatient mental health and emergency department (ED). Survey data were collected from all staff in the above roles who had worked for at least 6 months on one of the four participating clinical units. The exclusion criteria included anyone in a leadership role (eg, nurse manager) or anyone who was not in direct contact with patients (eg,

clerical staff responsible for administrative duties such as booking appointments for a nurse manager).

Survey data were collected between 30 September 2015 and 1 February 2016. During that time, the lead author visited each of the four participating units several times in order to recruit as many eligible full-time, part-time and casual front-line clinical staff as possible to complete a study survey. Non-probability convenience and snowball sampling procedures were used as it was not feasible to acquire accurate staffing numbers from the hospital/unit manager since casual staff were supplied by staffing agencies and assigned to a unit based on need. The on-site visits were spread across both the day shift and night shift so the researcher could meet and give surveys to as many eligible staff as possible. During each unit visit, a short oral presentation on the study's purpose, inclusion/exclusion criteria, survey characteristics (eg, voluntary, anonymous, cross sectional), and so on was used to solicit staff participation. Surveys were handed out to only those clinical staff who acknowledged that they met the study's inclusion criteria and were willing to participate in the study (four staff refused to take a survey). Respondents were asked to indicate the clinical unit they worked on; however, no individual identifiers were solicited (ie, survey data were anonymous). A drop box was placed on each participating unit to collect completed surveys. As a small inducement to participate, a \$20 gift card raffle draw was held on the final day of data collection on each unit. A returned completed survey by a respondent was seen as his/her consent to participate in the study.

Measures

Explanatory variables

The current study adhered to Zohar's conceptualisation of safety climate as employee's perceptions of management commitment to safety at two levels—senior and supervisory.^{11–13} The multilevel safety climate—that is, senior leadership support for safety and supervisory leadership support for safety—was measured using the Canadian Patient Safety Climate Survey (Can-PSCS).⁹ The Can-PSCS is a theory-based instrument that has strong psychometric properties validated by confirmatory factor analysis and is currently being used in health settings as part of Accreditation Canada's Qmentum Accreditation Program. The senior leadership support for safety scale has four items (eg, '*senior management considers patient safety when program changes are discussed*') and reflects staff perceptions of senior leadership commitment to PS. The supervisory leadership support for safety scale has two items (eg, '*my supervisor/manager seriously considers staff suggestions for improving patient safety*') and reflects staff perceptions of frontline-level leadership commitment to PS. Both scales were previously shown to have strong internal consistency reliability, $\alpha>0.80$.⁹

Staff perceptions of quality of teamwork on their respective units were measured using the Safety Attitudes Questionnaire teamwork climate scale. The scale has six items (eg, '*the physicians and nurses here work together as a*

well-coordinated team') and was previously shown to have good psychometric properties (eg, $\alpha=0.78$) in acute care settings.³¹

Outcome variable

The overall perceptions of PS scale was taken from the Agency for Healthcare Research and Quality Hospital Survey on Patient Safety Culture (HSOPSC).³² The HSOPSC overall perceptions of PS scale has four items (eg, '*we have patient safety problems in this unit*') and was previously shown to have strong internal consistency reliability, $\alpha>0.76$.³³

The senior leadership support for safety, supervisory leadership support for safety, teamwork climate and overall perceptions of PS were all measured using a 5-point agreement Likert response scale (1='strongly disagree' to 5='strongly agree'). All negatively phrased items were reverse coded so that a high score on an item corresponded to a high score on a scale. For each scale, a mean score was calculated if a respondent answered more than half of the questions in the scale.

Demographic variables

Clinical unit, tenure, gender and profession were all measured as categorical variables. Clinical unit contained four categories: general medicine, ICU, ED and mental health; unit tenure included three categories: 6–24 months, 2–5 years and >5 years. There were three categories for profession: clerical staff, nurses and AHPs, while gender contained two categories: male and female. See supplementary file 1 for the survey instrument used in the current study.

Analysis

All analyses were carried out using SPSS, version 22. Manual double entry of survey data was used to minimise data entry errors.³⁴ Cronbach's alpha values were calculated for senior leadership, supervisory leadership, teamwork climate and overall perceptions of PS to assess the reliability of these scales in the current data set.^{35 36} Demographic variables were transformed into multiple categorical variables (dummy variables) to use them in regression analyses. The categories of general medicine, 6–24 months, male and AHPs were used as reference groups.

Simple bivariate analyses (Pearson r) were carried out to assess the strength and significance of relationships among the dependent and non-demographic independent variables. Pearson r values were also examined to assess multicollinearity. High multicollinearity (ie, $r>0.8$) renders the results of regression analysis uninterpretable as it leads to statistically non-significant β coefficients even when R^2 is high and statistically significant.^{36 37} The residual scatter and probability-probability plots for overall perceptions of PS were examined to ensure that assumptions of multiple linear regression were met.^{35 36}

In order to test our study hypotheses, hierarchical regression analyses were used. Hierarchical regression

Table 1 Survey response rate by clinical unit

	Distributed	Refused survey at handout	Excluded (ineligible)	Returned	Response rate=returned ÷ (distributed+refused+ineligible)
Intensive care unit	66	2	0	49	49/68=72%
General medicine	49	0	0	45	45/49=92%
Emergency department	88	1	1	60	59/88=67%
Mental health	42	1	1	31	30/42=71%
Total	245	4	2	185	183/247=74%

analysis permits a researcher to examine the unique variance accounted for by a predictor, over and above the variance contributed by independent variables entered earlier in an analysis.³⁸ Demographic variables are typically good candidates for the first step in a hierarchical regression analysis³⁹ as they are static variables and should be entered in an analysis before the dynamic variables.³⁸ Hence, unit affiliation, and then tenure, and profession dummy variables were entered in block 1 and block 2 of the hierarchical regression analysis, respectively. The three predictors (ie, senior leadership support for safety, supervisory leadership support for safety and teamwork climate) were entered in block 3, and their associated interactions were entered in block 4. All predictors with interactions were centred in order to avoid problems of multicollinearity.³⁷ Significant interactions were plotted.

RESULTS

Response rate and sample characteristics

Table 1 shows the survey response rate for the current study. A total of 245 surveys were distributed. Four eligible clinical staff who refused to take a survey were added to the denominator for purposes of calculating the survey response rate and two returned surveys from respondents who indicated they had worked for <6 months on their clinical unit were excluded yielding a final denominator of 247.

The overall survey response rate was 74% (183/247). Response rates varied by unit (ED=67%; general medicine=92%, see **table 1**). It is possible that the 92% survey response rate on general medicine was facilitated by the physical space constraints of the unit—that is, presence of semiprivate patient rooms necessitated the charge nurse/unit clerk to ask all the staff to gather for a quick huddle when the primary researcher was on-site. These huddles made it easier for the researcher to build good rapport with staff and provided participants with an opportunity to complete the survey on the spot. Staff huddles were also conducted at other clinical units to help facilitate data collection but these occurred less frequently than on the general medicine unit.

Most study participants were female (89.6%), nurses (79.8%) and had a tenure of greater than 5 years on the unit (54.1%). The proportion of nurses (79.8%), AHPs (9.8%) and clerical staff (7.7%) in our survey respondents was similar to their proportion in participating

units' full-time staff where 82.5% were nurses, 9.7% were AHPs and 7.8% were clerics—see **table 2**. Other demographic information for participating clinical units' full-time nurses, AHPs and clerical staff was not available.

Bivariate analyses

Table 3 shows the results of the bivariate analyses and reveals significant relationships among the predictor and outcome variables. Multicollinearity was not a concern since all of the study's independent variables were correlated with each other at <0.6 and the variance inflation factor score for each of the three independent variables was <4³⁶. The Cronbach's α value for each study scale was >0.7 and values are shown in diagonal in **table 3**.

Hierarchical linear regression analyses

Table 4 shows the results of the hierarchical regression analyses. The clinical unit demographic variables, when entered in block 1 of the regression model, explained

Table 2 Demographic information of the whole sample (n=183)

	Frequency	%
Tenure		
6–24 months	24	13.1
2–5 years	51	27.9
>5 years	99	54.1
No response	9	4.9
Total	183	100
Gender		
Female	164	89.6
Male	16	8.7
No response	3	1.6
Total	183	100
Profession*		
Nurses	146	79.8
Allied health professional (AHP)	18	9.8
Clerical staff	14	7.7
No response	5	—
Total	183	100

*Professional breakdown of full-time staff reported in italics.

Table 3 Means, SDs and Pearson r correlations (n=183)

	Mean	SD	1	2	3	4
1. Senior leadership support for safety	3.01	0.94	0.87			
2. Supervisory leadership support for safety	3.61	1.02	0.490**	0.82		
3. Teamwork climate	3.61	0.67	0.402**	0.593**	0.78	
4. Overall PS perceptions	2.83	0.87	0.519**	0.408**	0.518**	0.81

**P<0.01.

PS, patient safety.

18% of the variance in overall perceptions of PS (adjusted $R^2=0.177$, p<0.001). The β coefficients for ICU ($\beta=0.548$, p<0.001) and mental health ($\beta=-0.591$, p<0.01) in model 1 were significant. These β coefficient values indicate that overall perceptions of PS scores were higher for ICU and lower for mental health compared with the reference category, general medicine. The staff demographic variables, when entered in block 2 of the regression model, did not explain a significant amount of variance in overall perceptions of PS (block 2 $\Delta R^2=0.036$, ns in **table 4**).

Senior leadership support for safety, supervisory leadership support for safety and teamwork climate when entered in block 3 of the regression model explained 24% of variance in overall perceptions of PS (block 3 $\Delta R^2=0.244$, p<0.001), over and above unit and demographic variables entered in previous blocks. β coefficients for senior

leadership support for safety ($\beta=0.360$, p<0.001) and teamwork climate ($\beta=0.420$, p<0.001) were significant.

Finally, the three interactions, when entered in block 4 of the regression model, did not explain a significant amount of additional variance in overall perceptions of PS (block 4 $\Delta R^2=0.015$, ns in **table 4**). However, the β coefficient of senior leadership and supervisory leadership interaction ($\beta=-0.135$, p<0.05) was significant. The significant interaction between senior and supervisory leadership is plotted in **figure 1**. **Figure 1** shows the relationship between perceptions of supervisory leadership for safety and overall PS when perceptions of senior leadership support for safety are high and when they are low. It shows that when perceptions of senior leadership support for safety are high, perceptions of supervisory leadership

Table 4 Results of hierarchical regression analysis (DV=overall perceptions of PS)

	Model 1, β	Model 2, β	Model 3, β	Model 4, β
Block 1—Unit affiliation				
ICU	0.548***	0.666***	0.485**	0.485**
ED	-0.132	-0.102	-0.145	-0.166
Mental health	-0.591**	-0.565**	-0.341*	-0.342*
Block 2—Staff demographics				
Tenure (2–5 years)	-0.068	0.114	0.095	
Tenure (>5 years)	-0.312	-0.099	-0.089	
Female	-0.299	-0.095	-0.126	
Nurses	-0.061	0.013	0.057	
Clerical staff	0.204	0.022	0.078	
Block 3—Predictor variables				
Senior leadership		0.360***	0.363***	
Supervisory leadership		-0.091	-0.151*	
Teamwork		0.420***	0.425***	
Block 4—Interactions				
Senior × supervisory				-0.135*
Senior × teamwork				0.122
Supervisory × teamwork				-0.029
Total R^2 (adjusted)	0.177***	0.189	0.435***	0.440
Change in R^2	0.192***	0.036	0.244***	0.015

Reference groups: general medicine, tenure (6–24 months), male and allied health professionals.

***P<0.001; **P<0.01; *P<0.05 (n=165).

DV, dependent variable; ED, emergency department; ICU, intensive care unit; PS, patient safety.




Figure 1 Interaction of senior leadership and supervisory leadership as predictors of overall perceptions of patient safety.

are not associated with overall PS perceptions; however, when perceptions of senior leadership support for safety are low, supervisory leadership becomes an important predictor of overall perceptions of PS. In total, the regression model accounted for 44% of the variance in overall perceptions of PS.

DISCUSSION

The survey results only partially supported hypothesis 1. Staff perceptions of senior leadership support for safety were shown to be significantly associated with overall perceptions of PS. However, the direct relationship between supervisory leadership support for safety and overall perceptions of PS was found to be non-significant. Other literature (reviewed above) suggests there is emerging empirical evidence of the positive impact of supportive supervisors on patient^{40 41} and staff⁴² outcomes. The survey we used solicited staff perceptions of only two *proactive* safety behaviours of a supervisor: (a) encouragement of clinical staff to follow established PS procedures and (b) consideration of staff suggestions for improving PS. It is possible, even likely, that clinical staff perceive safety related responsibilities of a supervisor much more broadly—for example, others have suggested that the ability to provide timely feedback for reported errors is seen as a central aspect of supervisory leadership support for safety.⁴³ Future research that operationalises supervisory leadership for safety in a broader way may reveal that this variable has a more pronounced effect on safety outcomes.

While we did not find evidence of a direct effect of supervisory leadership, our results highlight the importance of examining interactions and make a novel and

important contribution to the literature by showing a significant interaction between senior and supervisory leadership for safety. Our results suggest a compensatory effect where supervisory leaders' impact on overall perceptions of PS becomes strong and significant when front-line staff perceive poorer senior leadership support for safety (see *table 4* and *figure 1*). We found only one previous study that has empirically examined how senior leadership and supervisory leadership interact to explain variance in safety outcomes.²³ Recent qualitative work suggests a similar compensatory leadership result between unit and facility leaders⁴⁴ and between clinical and administrative leaders.⁴⁵ As noted, it is possible that a broader operational definition of supervisory leadership would reveal an even more pronounced compensatory effect. This line of enquiry requires further research and is especially relevant for loosely coupled organisations such as hospitals where front-line managers often hold considerable leeway while implementing policies created by senior leaders.¹⁴

Hypothesis 2 was also partially supported by the survey results as teamwork climate was shown to be significantly associated with overall perceptions of PS. These results are consistent with past empirical research that has shown staff perceptions of teamwork climate significantly impact patient outcomes—for example, adverse events, mortality rates^{22 28 29} and staff outcomes (eg, burnout).²⁷

Our results showed that, compared with being on a general medicine unit, working in ICU was associated with more positive perceptions of overall PS, and working on a mental health unit was associated with poorer perceptions of overall PS. These results are consistent with previous research showing that staff perceptions of domain-specific climates differ between hospital work areas.⁴⁶ ICUs may attract more resources due to greater case acuity, specialised focus and higher status of physicians and nurses. In contrast, clinical areas such as mental health have been noted for resource scarcity, quality issues and inadequate staff training⁴⁷—all factors that may lead to negative perceptions of PS. Roots of variation in safety climate across clinical areas should be the subject of further inquiry. Our leadership results may suggest a starting point for practices to raise the safety climate levels in areas such as mental health.

Limitations and future research

This study was cross sectional and therefore causal associations between predictors and outcome cannot be established. Also, self-reported measures were used that are subject to social desirability biases.⁴⁸ However, assuring survey participants' anonymity, as was done in the current study, can minimise socially desirable responding.⁴⁹

Physicians were not included in the current study since only a small number of full-time physicians worked on the participating general medicine and mental health units and physicians are often not physically present on a clinical unit throughout a shift making their recruitment using the study's data collection procedures difficult.

Physicians are also more likely to be informally seen as team leaders by other clinical staff and the current study did not include clinicians in leadership roles.

The current study's predictor and outcome variables were taken from the same survey. Hence, common methods variance is likely to inflate the magnitude of the relationships we examined. Finally, convenience and snowball sampling procedures were used and data come from a single large community hospital. It is recommended that future research test the validity of the current study's inferences in other types of clinical units (eg, surgery or paediatrics), professions (eg, physicians) and hospitals (eg, small community or teaching) using larger multisite samples.

Implications for practice

The relational practices—for example, providing vision, support and constructive feedback—of formal healthcare leaders have been found to be associated with better staff and patient outcomes.^{20 41} While healthcare institutions should focus on recruiting and retaining individuals possessing relational competencies into leadership roles, healthcare leaders must work hard to prioritise safety over efficiency and be cognisant of the limitations of certain performance indicators—for example, average length of stay, average cost per discharge—which emphasise efficiency over safety. In settings where leadership support for safety is lagging, attention can be directed to a small but growing evidence base that suggests leadership for quality and safety can be built as part of interventions to improve care. Organisations and health systems are encouraged to view leadership for safety as a modifiable element that can be fostered rather than a fixed aspect of context that is either present or absent.⁴⁴

Health organisations are under increasing pressure to simultaneously contain costs and preserve quality of care. These circumstances necessitate higher levels of collaboration and teamwork among healthcare professionals. At the same time, empirical evidence suggests that clinicians belonging to different professions—for example, physicians, nurses—hold different perspectives on and expectations for collaborative behaviours at the frontlines.⁵⁰ Therefore, healthcare organisations must provide on-site interprofessional collaborative workshops on topics that can strengthen working relationships including conflict management, negotiation skills and stress management.⁵¹

The adult inpatient mental health staff held poorer perceptions of overall PS compared with other clinical units. Mental health units may experience greater threats to staff safety as a result of social and clinical conditions particular to patients on those units.⁵² There is evidence to suggest that staff on mental health units experience three times as much job-related violence compared with their peers on non-mental health units.⁵³ Our results, along with other empirical research demonstrating resource scarcity and increased risk of violence in these settings, strongly suggest that healthcare managers/leaders must devote more attention and perhaps resources to patient and staff safety on these units.

CONCLUSION

The '*To Err is Human*'²⁶ report highlighted serious quality and safety deficiencies in healthcare delivery systems and in doing so energised healthcare professionals/researchers to design, evaluate and implement safety improvement strategies at the frontlines. Indeed, implementation of standardised clinical interventions—for example, hand hygiene guidelines³ and surgical checklists⁴—has reduced preventable errors and associated patient harm. There is also increasing evidence of the influence of contextual factors on quality of care and safety outcomes^{5 27}; however, certain literature gaps still remain—for example, an imprecise conceptualisation of multilevel safety climate and a dearth of empirical research on its interactive effect on safety outcomes. The results of the current study suggest that senior leaders' support for safety and teamwork climate significantly impact nursing, allied health and clerical staff perceptions of PS in an acute care setting. However, when senior leaders provide insufficient support for safety, a supervisor is still able to positively mould front-line staff overall perceptions of PS. This particular finding is noteworthy as it highlights the underexplored compensatory nature of multilevel safety climate where leadership support at one level can counterbalance a lack of leadership support at another level. Our results begin to address some of the organisational safety literature gaps and suggest some contextual aspects of organisations that could benefit from increased attention in PS science.

Acknowledgements We thank Dr Dennis Raphael, Dr Whitney Berta and Dr Mary Fox for their contributions as members of the oral defence committee for SZ's PhD dissertation. Most importantly, we acknowledge and thank those in the hospital where this study took place including the Director of Medical Education, Dr Zaev Wulffhart, as well as other senior leaders for their continuous support, the front-line managers on each of the study units for facilitating access to their units, and all of the staff who generously agreed to participate in this study.

Contributors SZ designed the study, collected and analysed the data, and drafted and revised the paper. LRG, HJW and KT contributed to overall study design, advised on data collection and analysis, and revised and approved the manuscript. LB contributed to study design, advised on data collection and facilitated access to the research site.

Competing interests None declared.

Ethics approval Ethics approval was obtained from the participating hospital's ethics board and the Human Participants Review Sub-Committee of York University's Ethics Review Board (certificate number STU 2016-016).

Provenance and peer review Not commissioned; externally peer reviewed.

Data sharing statement The data set used during the current study is available from the corresponding author on reasonable request.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

REFERENCES

1. Baker GR, Norton PG, Flintoft V, et al. The Canadian adverse events study: the incidence of adverse events among hospital patients in Canada. *CMAJ* 2004;170:1678–86.

2. Etchells E, Mittmann N, Koo M, et al. The economics of patient safety in acute care: technical report. Canadian Patient Safety Institute. 2012. <http://www.patientsafetyinstitute.ca/en/toolsResources/Research/commissionedResearch/EconomicsofPatientSafety/Documents/Economics%20of%20Patient%20Safety%20-%20Acute%20Care%20-%20Final%20Report.pdf> (Accessed 12 May 2018).
3. Larson E, Goldmann D, Pearson M, et al. Measuring hand hygiene adherence: overcoming the challenges. The Joint Commission. 2009 https://www.jointcommission.org/assets/1/18/lh_monograph.pdf (Accessed 12 May 2018).
4. World Health Organization & WHO Patient Safety. *Implementation manual: WHO surgical safety checklist (first edition)*. Geneva: World Health Organization, 2008.
5. Bosk CL, Dixon-Woods M, Goeschel CA, et al. Reality check for checklists. *Lancet* 2009;374:444–5.
6. Pronovost P, Needham D, Berenholtz S, et al. An intervention to decrease catheter-related bloodstream infections in the ICU. *N Engl J Med* 2006;355:2725–32.
7. Weick KE, Sutcliffe KM. *Managing the Unexpected*. 2nd edn. San Francisco, CA: John Wiley & Sons, Inc, 2007.
8. Vogus TJ, Sutcliffe KM, Weick KE. Doing no harm: enabling, enacting, and elaborating a culture of safety in health care. *Acad Manag Perspect* 2010;24:60–77.
9. Ginsburg LR, Tregunno D, Norton PG, et al. 'Not another safety culture survey': using the Canadian patient safety climate survey (Can-PSCS) to measure provider perceptions of PSC across health settings. *BMJ Qual Saf* 2014;23:162–70.
10. Singer SJ, Vogus TJ. Safety climate research: taking stock and looking forward. *BMJ Qual Saf* 2013;22:1–4.
11. Zohar D. Safety climate in industrial organizations: theoretical and applied implications. *J Appl Psychol* 1980;65:96–102.
12. Zaheer S, Ginsburg L, Chuang YT, et al. Patient safety climate (PSC) perceptions of frontline staff in acute care hospitals: examining the role of ease of reporting, unit norms of openness, and participative leadership. *Health Care Manage Rev* 2015;40:13–23.
13. Zohar D. Safety climate and beyond: a multi-level multi-climate framework. *Saf Sci* 2008;46:376–87.
14. Zohar D, Hofmann DA. Organizational culture and climate. In: Kozlowski SWJ, ed. *The Oxford handbook of organizational psychology*. New York: Oxford University Press, Inc, 2012:643–66.
15. Yun S, Faraj S, Sims HP. Contingent leadership and effectiveness of trauma resuscitation teams. *J Appl Psychol* 2005;90:1288–96.
16. Frankel A, Grillo SP, Pittman M, et al. Revealing and resolving patient safety defects: the impact of leadership WalkRounds on frontline caregiver assessments of patient safety. *Health Serv Res* 2008;43:2050–66.
17. Thomas EJ, Sexton JB, Neilands TB, et al. The effect of executive walk rounds on nurse safety climate attitudes: a randomized trial of clinical units [ISRCTN85147255] [corrected]. *BMC Health Serv Res* 2005;5:28.
18. Pronovost PJ, Weast B, Bishop K, et al. Senior executive adopt-a-work unit: a model for safety improvement. *Jt Comm J Qual Saf* 2004;30:59–68.
19. Tucker AL, Singer SJ, Hayes JE, et al. Front-line staff perspectives on opportunities for improving the safety and efficiency of hospital work systems. *Health Serv Res* 2008;43:1807–29.
20. Spence Laschinger HK, Leiter MP. The impact of nursing work environments on patient safety outcomes. *JONA: The Journal of Nursing Administration* 2006;36:259–67.
21. Cole E, Crichton N. The culture of a trauma team in relation to human factors. *J Clin Nurs* 2006;15:1257–66.
22. Mardon RE, Khanna K, Sorra J, et al. Exploring relationships between hospital patient safety culture and adverse events. *J Patient Saf* 2010;6:226–32.
23. Zohar D, Livne Y, Tenne-Gazit O, et al. Healthcare climate: a framework for measuring and improving patient safety. *Crit Care Med* 2007;35:1312–7.
24. Sutcliffe KM, Lewton E, Rosenthal MM. Communication failures: an insidious contributor to medical mishaps. *Acad Med* 2004;79:186–94.
25. Berlin JM, Carlström ED. The 20-minute team--a critical case study from the emergency room. *J Eval Clin Pract* 2008;14:569–76.
26. Kohn LT, Corrigan JM, Donaldson MS. *To err is human: building a safer health system*. Washington, DC: National Academy Press, 2000.
27. Bowers L, Nijman H, Simpson A, et al. The relationship between leadership, teamworking, structure, burnout and attitude to patients on acute psychiatric wards. *Soc Psychiatry Psychiatr Epidemiol* 2011;46:143–8.
28. Mazzocco K, Petitti DB, Fong KT, et al. Surgical team behaviors and patient outcomes. *Am J Surg* 2009;197:678–85.
29. Buist MD, Moore GE, Bernard SA, et al. Effects of a medical emergency team on reduction of incidence of and mortality from unexpected cardiac arrests in hospital: preliminary study. *BMJ* 2002;324:387–90.
30. Singer SJ, Vogus TJ. Reducing hospital errors: interventions that build safety culture. *Annu Rev Public Health* 2013;34:373–96.
31. Sexton JB, Holzmueller CG, Pronovost PJ, et al. Variation in caregiver perceptions of teamwork climate in labor and delivery units. *J Perinatol* 2006;26:463–70.
32. AHRQ. Hospital survey on patient safety culture. 2015. <http://www.ahrq.gov/professionals/quality-patient-safety/patientsafetyculture/hospital/> (Accessed 13 May 2018).
33. Etchegaray JM, Thomas EJ. Comparing two safety culture surveys: safety attitudes questionnaire and hospital survey on patient safety. *BMJ Qual Saf* 2012;21:490–8.
34. Paulsen A, Overgaard S, Lauritsen JM. Quality of data entry using single entry, double entry and automated forms processing—an example based on a study of patient-reported outcomes. *PLoS One* 2012;7:e35087.
35. Dancey CP, Reidy JG, Rowe R. *Statistics for the health sciences: a non-mathematical introduction*. London: SAGE Publications, Inc, 2012.
36. Katz MH. *Multivariable analysis: a practical guide for clinicians*. 2nd edn. Cambridge: Cambridge University Press, 2006.
37. Tabachnick BG, Fidell LS. *Using multivariate statistics*. 6th edn. London: Pearson Education, Inc, 2013.
38. Petrocelli JV. Hierarchical multiple regression in counseling research: common problems and possible remedies. *Meas Eval Couns Dev* 2003;39:29–2.
39. Cohen J, Cohen P. *Applied multiple regression/correlation analysis for the behavioral sciences*. 2nd edn. Hillsdale, NJ: Erlbaum, 1983.
40. Brady Germain P, Cummings GG. The influence of nursing leadership on nurse performance: a systematic literature review. *J Nurs Manag* 2010;18:425–39.
41. Wong CA, Cummings GG, Ducharme L. The relationship between nursing leadership and patient outcomes: a systematic review update. *J Nurs Manag* 2013;21:709–24.
42. Galletta M, Portoghesi I, Battistelli A, et al. The roles of unit leadership and nurse-physician collaboration on nursing turnover intention. *J Adv Nurs* 2013;69:1771–84.
43. Evans SM, Berry JG, Smith BJ, et al. Attitudes and barriers to incident reporting: a collaborative hospital study. *Qual Saf Health Care* 2006;15:39–43.
44. Ginsburg L, Easterbrook A, Berta W, et al. Implementing frontline worker-led quality improvement in nursing homes: getting to "how". *The Joint Commission Journal on Quality and Patient Safety* 2018;44:526–35.
45. van der Zijpp TJ, Niessen T, Eldh AC, et al. A bridge over turbulent waters: illustrating the interaction between managerial leaders and facilitators when implementing research evidence. *Worldviews Evid Based Nurs* 2016;13:25–31.
46. Singer SJ, Gaba DM, Falwell A, et al. Patient safety climate in 92 US hospitals: differences by work area and discipline. *Med Care* 2009;47:23–31.
47. Armstrong P, Banerjee A, Szebehely M, et al. *They deserve better: the long-term care experience in Canada and Scandinavia*. Ottawa: Canadian Center for Policy Alternatives, 2009.
48. Foley KL, Manuel J, Vitalins M. The utility of self-report in medical outcomes research. *Evidence-based Healthcare and Public Health* 2005;9:263–4.
49. Randall DM, Fernandes MF. The social desirability response bias in ethics research. *Journal of Business Ethics* 1991;10:805–17.
50. Copnell B, Johnston L, Harrison D, et al. Doctors' and nurses' perceptions of interdisciplinary collaboration in the NICU, and the impact of a neonatal nurse practitioner model of practice. *J Clin Nurs* 2004;13:105–13.
51. Nair DM, Fitzpatrick JJ, McNulty R, et al. Frequency of nurse-physician collaborative behaviors in an acute care hospital. *J Interprof Care* 2012;26:115–20.
52. Anderson A, West SG. Violence against mental health professionals: when the treater becomes the victim. *Innov Clin Neurosci* 2011;8:34–9.
53. Friedman RA. Violence and mental illness--how strong is the link? *N Engl J Med* 2006;355:2064–6.