Prevalence and clinical characteristics of inappropriate myocardial perfusion imaging tests at a community hospital

Nadia Isabel Abelhad, Kendall J Kiser, Andres Hughes, Michael A Hust, Enrique Garcia Sayan, Siddharth K Prakash

To cite: Abelhad NI, Kiser KJ, Hughes A, *et al.* Prevalence and clinical characteristics of inappropriate myocardial perfusion imaging tests at a community hospital. *BMJ Open Quality* 2019;**8**:e000487. doi:10.1136/bmjopen-2018-000487

Received 14 August 2018
Revised 4 April 2019
Accepted 27 April 2019

© Author(s) (or their employer(s)) 2019. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

Department of Internal Medicine, University of Texas Health Science Center at Houston, Houston, Texas, USA

Correspondence to

Dr Siddharth K Prakash; siddharth.k.prakash@uth.tmc.edu

BACKGROUND

Myocardial perfusion imaging (MPI) is a cornerstone for the diagnosis and management of patients with coronary artery disease (CAD). However, the persistence of inappropriate MPI tests as documented by meta-analyses undermines their clinical utility.¹ Inappropriate MPI tests reduce diagnostic accuracy and increase false-positive results that can subject patients to unnecessary invasive procedures with additional risks, including radiation exposure, renal insufficiency and bleeding.^{2 3} Indirect effects at our hospital include prolongation of waiting time for procedures and referrals (currently 5 months for outpatient MPI) and consumption of limited healthcare resources (\$350–\$500 per test)⁴ in a predominately uninsured population.⁵ Inappropriate tests led to more than \$500 million of additional costs in 2010 alone.⁶

To address these challenges, cardiovascular societies promulgated appropriate use criteria (AUC) for MPI, and Congress directed the Centers for Medicare & Medicaid Services to consider AUC in reimbursement decisions.⁷ Risk calculators that emerged from these efforts are useful to identify appropriate patients who may benefit from MPI tests but remain underused. To investigate the utility of this approach, we evaluated the prevalence and characteristics of inappropriate MPI orders at one general community hospital.

METHODS

We conducted a 3-month retrospective electronic medical record review of all patients (n=324) at a single hospital who were referred for MPI within 12 months of coronary angiography. Subjects who did not undergo prior MPI and subjects without complete clinical information were excluded (n=262). The appropriateness of MPI orders was adjudicated using the 2009 criteria for radionuclide

imaging.¹ Pretest probabilities of obstructive CAD were estimated using revised Diamond-Forrester Index (DFI) and atherosclerotic cardiovascular disease (ASCVD) calculators. Positive predictive values (PPVs) were calculated for >50% angiographic stenosis. Comparisons of categorical variables were evaluated using χ^2 or Fisher exact tests. The severity of perfusion defects was compared using Spearman's rank-order tests.

RESULTS

Sixty-five subjects (49% female, 58% current or former smokers) met all inclusion criteria. The distribution of appropriate (78%), inappropriate (17%) and indeterminate (5%) MPIs was similar to historical data. However, the normalcy rate (30%) was significantly less than expected (90%) and was correlated with the prevalence of obesity. Elevated body mass index (34.0 in men and 37.1 in women) may result in decreased normalcy rates due to breast and/or abdominal attenuation.⁸ Seventy percent of MPI referrals, including all inappropriate referrals, originated from outpatient clinics. In contrast, none of the hospital inpatient (22%) or emergency department (8%) referrals were inappropriate. Inappropriate referrals were most frequently requested for patients with low pretest probabilities of CAD, as defined by a DFI of <10% (74.1%) or an ASCVD of <5% (40.7%). Inappropriately referred patients were more likely to present with preserved systolic function and were less likely to present with angina. The most common presenting symptom of appropriately referred patients was atypical angina (51.9%). The mean HbA1c values (5.9% vs 7.4%, p=0.02) and estimated ASCVD 10-year risk (9.0% vs 17.4%, p=0.08) of inappropriately referred patients were lower than those for appropriately referred patients (table 1). The PPV of

Table 1 Comparison of patients who underwent appropriate and inappropriate myocardial perfusion imaging stratified by pretest probabilities

Characteristic	Inappropriate (n=10, 17%)		Appropriate (n=49, 78%)	
	Mean	σ	Mean	σ
Ejection fraction	58%	4.5	48%	14.7
Referral location	Outpatient	100%	Inpatient	100%
DFI*	<10%	0.5	>10%–90%	0.7
ASCVD score†	9.0%	8.0	17.4%	15.4
A1c level‡	5.9%	0.8	7.4%	2.0

*P=0.06.

†P=0.08.

‡P=0.02.

ASCVD, atherosclerotic cardiovascular disease; A1c, haemoglobin A1c; DFI, Diamond-Forrester Index.

inappropriate MPI studies (22%) was also lower than that of appropriate studies (49%, $p=0.07$; table 2).

CONCLUSION

In a contemporary hospital-based cohort with prevalent CAD, we found that inappropriate MPI orders remain common and are related to predictable factors. Most inappropriate referrals originated from a small number of providers and involved low-probability patients who had normal left ventricular systolic function and did not present with typical anginal chest pain. The lower PPVs of inappropriate studies, in comparison with those of appropriate studies, are associated with diminished clinical utility for diagnosis and treatment.⁹ Inappropriate studies also increase the likelihood of potential harm from unnecessary interventions and exacerbate the misdirection of limited clinical resources. These observations highlight the potential financial and clinical impacts of educational interventions or interactive pop-up alerts in the electronic medical record to reduce inappropriate MPI referrals. We are currently assessing the effectiveness

Table 2 The PPV of inappropriate MPI tests was lower than that of appropriate MPI tests

Test result	Inappropriate (n=10)		Appropriate (n=52)	
	+CAD	–CAD	+CAD	–CAD
Abnormal MPI	2	7	19	20
Normal MPI	0	1	3	10
PPV	22%		49%	

CAD, at least one >50% coronary lesion by coronary angiography; MPI, myocardial perfusion imaging; PPV, positive predictive value.

of targeted information sessions at the same hospital to promote provider awareness and to change test ordering patterns.¹⁰

Acknowledgements We sincerely thank LBJ Hospital, the Harris Health IT department for compiling our list of patients, the cardiology staff at LBJ for imaging and procedural reports, and the Harris Health Quality and Safety Committee for permitting our research.

Contributors SP and ES supervised the study. NA, KK, MH and AH abstracted clinical data. NA wrote the manuscript. SP, ES, KK and NA edited the manuscript.

Funding SP was supported by the Cheves and Isabella Smythe Distinguished Professorship in Internal Medicine.

Competing interests None declared.

Patient consent for publication Not required.

Ethics approval The Committee for the Protection of Human Subjects at McGovern Medical School and the Harris Health System Office of Research and Sponsored Programs reviewed and approved the study protocol.

Provenance and peer review Not commissioned; externally peer reviewed.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

REFERENCES

1. Hendel RC, Berman DS, Di Carli MF, *et al.* ACCF/ASNC/ACR/AHA/ASE/SCCT/SCMR/SNM 2009 appropriate use criteria for cardiac radionuclide imaging: a report of the American College of cardiology Foundation appropriate use criteria Task force, the American Society of nuclear cardiology, the American College of radiology, the American heart association, the American Society of echocardiography, the Society of cardiovascular computed tomography, the Society for cardiovascular magnetic resonance, and the Society of nuclear medicine. *Circulation* 2009;119:e561–87.
2. Grayburn PA, Hillis LD. Cardiac events in patients undergoing noncardiac surgery: shifting the paradigm from noninvasive risk stratification to therapy. *Ann Intern Med* 2003;138:506–11.
3. Wijesundera DN, Beattie WS, Austin PC, *et al.* Non-Invasive cardiac stress testing before elective major non-cardiac surgery: population based cohort study. *BMJ* 2010;340:b5526.
4. U.S. Centers for Medicare and Medicaid Services. 2018 Physician Fee Schedule [Internet]. Baltimore, MD: U.S. Centers for Medicare & Medicaid Services, 2018. Available: <http://www.cms.gov/apps/physician-fee-schedule/search/search-results.aspx?Y=1&T=0&HT=1&CT=3&H1=78451&H2=78452&H3=78453&H4=78454&H5=78456&M=5> [Accessed 25 Feb 2019].
5. Census Bureau (US). United States Census Bureau QuickFacts: Harris County, Texas [Internet]. Washington, DC: U.S. Department of Commerce, 2018. Available: www.census.gov/quickfacts/fact/table/harriscountytexas/PST045217 [Accessed 29 Feb 2019].
6. . Report to the Congress: Medicare payment policy. Washington, D.C. Medicare Payment Advisory Commission; 2005.
7. Ladapo JA, Blecker S, O'Donnell M, *et al.* Appropriate use of cardiac stress testing with imaging: a systematic review and meta-analysis. *PLoS One* 2016;11:e0161153.
8. Elgendy IY, Mahmoud A, Shuster JJ, *et al.* Outcomes after inappropriate nuclear myocardial perfusion imaging: a meta-analysis. *J Nucl Cardiol* 2016;23:680–9.
9. Green R, Cantoni V, Petretta M, *et al.* Negative predictive value of stress myocardial perfusion imaging and coronary computed tomography angiography: a meta-analysis. *J Nucl Cardiol* 2018;25:1588–97.
10. Ladapo JA, Blecker S, Douglas PS. Physician decision making and trends in the use of cardiac stress testing in the United States: an analysis of repeated cross-sectional data. *Ann Intern Med* 2014;161:482–90.