Testing a proactive approach to frailty identification: the electronic frailty index

Nathan Devereux,¹ Graham Ellis,^{1,2} Laura Dobie,¹ Paul Baughan,^{1,3} Thomas Monaghan¹

To cite: Devereux N, Ellis G, Dobie L, *et al.* Testing a proactive approach to frailty identification: the electronic frailty index. *BMJ Open Quality* 2019;8:e000682. doi:10.1136/bmjopen-2019-000682

► Additional material is published online only. To view please visit the journal online (<http://dx.doi.org/10.1136/bmjopen-2019-000682>).

Received 5 March 2019
Revised 28 June 2019
Accepted 6 August 2019

INTRODUCTION

Frailty is the manifestation of ageing that is associated with poor outcomes, including increased risk of disability, hospital admission, institutional care or death.¹ It can have a considerable impact on a person's quality of life, and lead to increased use of primary care and unplanned secondary care services.

However, frailty is not an inevitable part of ageing, and an individual's frailty state is not constant: it can be improved or exacerbated.² Identifying people with frailty in primary care, and targeting them with appropriate interventions, can delay the progression of this condition.³

Aim

In order to improve outcomes for people with frailty, we needed to develop and test a consistent method to identify people, which could be used in a community setting and implemented at a national scale throughout Scotland.

METHODS AND RESULTS

Understand

We selected the electronic frailty index (eFI) as the method to explore in Scotland, as it had been validated in England against a population of 900 000 and uses existing data to identify a person's condition.¹ It was unclear whether it would accurately identify people with frailty using Scottish data and IT infrastructure.

Test and refine

We tested the eFI on a population of 75 311 (12 496 aged ≥65) across nine general practitioner (GP) practices in three health and social care partnerships in Scotland. Read code data from the participating GP practices was used to stratify the population into mild, moderate or severe frailty. More information

on eFI deficits and cut-offs can be found in the appendix.

We reviewed a sample of the eFI results for three practices, totalling 135 people. The GP in each practice compared the eFI result against the Rockwood clinical frailty scale⁴ and marked whether they agreed with the assessment or not. Table 1 compares our results with the external validation cohort in the original study.

We then refined the presentation of the results to focus on individuals experiencing the greatest change in their frailty status, in order to ensure that those with greatest need were identified for review by community teams. This ensured that the numbers of individuals identified for a frailty case review could be matched with the capacity of professionals to conduct assessments. The Scottish Clinical Information Management in Practice GP group independently reviewed read codes used in Scotland and added these to the tool to ensure that codes commonly used in Scotland were included under the relevant deficits.

Implement and spread

Following successful testing, we wanted to make this tool available to all GP practices in Scotland to identify people with frailty in their population who could benefit from community-based support.

We worked with Information Services Division to make the adapted eFI available through the Scottish Primary Care Information Resource (SPIRE).⁵ SPIRE is a portal that facilitates the management of information at a practice level.

As well as providing a list of people in the GP practice who are living with frailty, SPIRE can highlight the following groups of people:

- People who escalate to being moderately frail during the previous 6 months.

© Author(s) (or their employer(s)) 2019. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

¹Improvement Hub, Healthcare Improvement Scotland, Edinburgh, UK

²University Hospital Monklands, NHS Lanarkshire, Bothwell, UK

³Dollar Health Centre, NHS Forth Valley, Stirling, UK

Correspondence to

Laura Dobie;
laura.dobie@nhs.net

Table 1 Original eFI study external validation cohort results compared with our testing

	Original study external validation cohort (65–95)	Our testing (age 65+)
No. male	227 043	5460
No. female	288 964	7036
Male (%)	44	44
Female (%)	56	56
Mild frailty (%)	37	35
Moderate frailty (%)	16	15
Severe frailty (%)	4	5

- People who are moderately frail and have experienced the greatest change during a 6-month period.
- People who escalate to being severely frail during a 6-month period.

DISCUSSION

We tested and adapted a clinically validated tool to reflect the capacity of community-based teams. This adaptation was crucial to successful adoption of the tool by the test practices.

Working in partnership with colleagues in National Services Scotland, we have now made this tool available through infrastructure that enables its implementation at scale.

Our test practices were concentrated in urban areas and areas of greater deprivation. This may have skewed the eFI results, as the validation demonstrated higher eFI scores in areas with higher levels of deprivation.¹ However, Dollar Health Centre, a semi-rural GP practice with low levels of deprivation, is also seeing the benefit of using the eFI with their population. This practice uses the tool at their multidisciplinary team meeting to highlight people who may need additional support. This is often a referral to an anticipatory care planning (ACP) nurse, who will visit the patient at home, conduct a holistic assessment and initiate ACP discussions. The team expect that this will increase the number of completed ACPs and reduce hospital admissions, although at present there is insufficient data to demonstrate this.

The implementation of a robust frailty identification tool will enable teams, organisations and national programmes to focus preventative support on those people who are likely to benefit the most. This will improve the quality of life of people who become frail and reduce demand on unplanned services that are associated with crisis moments.

Acknowledgements Dr Andrew Clegg, University of Leeds, gave permission to use the eFI algorithm. Glasgow City, West Dunbartonshire and Midlothian Health and Social Care Partnerships participated in the eFI testing. Hugh O'Pray, NHS Greater Glasgow and Clyde, extracted GP read code data in NHS Greater Glasgow and Clyde. Mohini Padayachee, NHS Lothian, extracted GP read code data in NHS Lothian. National Services Scotland are making the eFI available on the Scottish Primary Care Information Resource. The Scottish Clinical Information Management in Practice GP group reviewed the GP read codes included in the eFI.

Contributors ND is the Improvement Advisor who led this improvement project. ND supported GP practices to test the eFI. GE provided clinical expertise to the project. LD provided communications support to the project and drafted the article. PB tested the eFI in his GP practice and provided information on how his practice is using the tool. TM is the Programme Team Lead and is the guarantor for the content of the paper. TM supported GP practices to test the eFI.

Funding The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

Competing interests None declared.

Patient consent for publication Not required.

Provenance and peer review Not commissioned; externally peer reviewed.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

REFERENCES

1. Clegg A, Bates C, Young J, *et al.* Development and validation of an electronic frailty index using routine primary care electronic health record data. *Age Ageing* 2016;45:353–60.
2. British Geriatrics Society. Fit for frailty 1: consensus best practice guidance for the care of older people living in community and outpatient settings, 2014. Available: https://www.bgs.org.uk/sites/default/files/content/resources/files/2018-05-23/fff_full.pdf [Accessed 4 Jun 2019].
3. Lee L, Heckman G, Molnar FJ. Frailty: identifying elderly patients at high risk of poor outcomes. *Can Fam Physician* 2015;61:227–31.
4. Rockwood K, Song X, MacKnight C, *et al.* A global clinical measure of fitness and frailty in elderly people. *CMAJ* 2005;173:489–95.
5. NHS National Services Scotland. Scottish primary care information resource. Available: <https://spire.scot/professional/> [Accessed 26 Feb 2019].