Evaluating the effect of nurse-initiated discussion of infection management during ICU bedside rounds

Linda Dresser,^{1,2} Madeleine S Stephen,³ Mark McIntyre,^{1,2} Linda Jorgoni,¹ Sarah C J Jorgensen,^{1,4} Sandra Nelson,⁵ Chaim Bell,^{1,6,7} Andrew M Morris^{1,7}

To cite: Dresser L, Stephen MS, McIntyre M, *et al.* Evaluating the effect of nurse-initiated discussion of infection management during ICU bedside rounds. *BMJ Open Quality* 2020;9:e001037. doi:10.1136/bmjopen-2020-001037

Received 29 May 2020

Revised 20 August 2020

Accepted 14 October 2020

© Author(s) (or their employer(s)) 2020. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

¹Antimicrobial Stewardship Program, Sinai Health/University Health Network, Toronto, Ontario, Canada

²Leslie Dan Faculty of Pharmacy, University of Toronto, Toronto, Ontario, Canada

³Faculty of Medicine, University of Ottawa, Ottawa, Ontario, Canada

⁴Department of Pharmacy, Sinai Health, Toronto, Ontario, Canada

⁵Department of Pharmacy, Sunnybrook Health Sciences Centre, Toronto, Ontario, Canada

⁶Division of Internal Medicine, Sinai Health, Toronto, Ontario, Canada

⁷Department of Medicine, University of Toronto, Toronto, Ontario, Canada

Correspondence to

Dr Linda Dresser;
Linda.dresser@utoronto.ca

INTRODUCTION

Intensive care unit (ICU) antimicrobial stewardship (AMS) interventions generally do not include nurses although their role is increasingly recognised.^{1–4} Previous literature described inclusion of nurses and nurse-led AMS interventions outside of ICU daily bedside rounds.^{5,6} After using a structured quality improvement (QI) evaluation of the daily bedside rounding model of the ICU clinical team we hypothesised the addition of ‘infection’ as an entity in the ICU nurses’ structured daily report would improve inter-professional discussion of infection management.

METHODS

Setting

Thirty-bed, tertiary-care, medical-surgical ICU at an academic centre with an established AMS programme.⁷

Design

We used the QI improvement methodology of Plan, Do, Study, Act cycle to frame this project.

Phase 1: Plan: We identified process gaps in infection management occurring during ICU interprofessional rounds.

Workflow analysis identified that the nursing daily report provided unstructured information about infection status. Infection management discussions were limited to ICU pharmacists and physicians.

Focus groups found that nurses frequently did not know the indication for prescribed antimicrobials but perceived this knowledge to be important in providing safe and effective care.

Therefore, we proposed the introduction of ‘infection’ as a separate entity in the daily nurse report to provide structure for essential information dissemination and allow routine nursing inclusion in team discussions

of infection management. Using the theoretical domains framework of behaviour change, we developed an intervention focused on nurses’ capability, motivation and opportunity to participate in discussion of infection management.^{8,9} Our phased approach used education, persuasion, training and enablement to facilitate behaviour change.

Phase 2: Do: We engaged ICU stakeholders, including nursing leaders, to design, implement and support inclusion of ‘infection’ in the nurse report.

During meetings with ICU physicians and pharmacists, we highlighted how prior AMS rounds had equipped them with a skill set to incorporate the framework for infection management (table 1) into their daily practice. They now needed to facilitate nurse participation.

Forums with ICU nurses provided the rationale for inclusion of ‘infection’ and addressed concerns regarding nursing role in management of infection. Similar forums with ICU pharmacists addressed concerns about redundancy of roles during bedside rounds.

We achieved consensus among all stakeholders regarding process and launch date.

At the launch, the AMS nurse provided ICU nurses with group or individual education. Stickers in the bedside nursing clinical summary tool (Kardex) served as visual reminders. The AMS nurse supported nurses by attending daily rounds and giving real-time feedback on their reporting.

Phase 3: Study: Post-implementation training and feedback to improve intervention acceptance and sustainability was performed.

Based on nursing feedback, we modified the training approach for new nurses, to include role-playing and scripts. Audit and feedback during rounds continued and additional visual cues were added to the ICU daily clinical flow sheet.

Table 1 Pre-intervention and post-intervention changes in inclusion of infection management decision-making framework components during daily bedside ICU rounds

Decision-making component	Pre-intervention n=95 n (%)	Post-intervention n=217 n (%)	P value
Evidence of infection	7 (7.4)	158 (72.8)	<0.001
Focus of infection	17 (17.9)	158 (72.8)	<0.001
Likely pathogens of infection	16 (16.8)	154 (71.0)	<0.001
Intention of therapy	16 (16.8)	165 (76.0)	<0.001
Current day of antimicrobial therapy	20 (21.1)	174 (80.2)	<0.001
Tailoring antimicrobial therapy	66 (69.5)	113 (52.1)	0.004
Expectations adjusted	22 (23.2)	117 (53.9)	<0.001
Planned duration	51 (53.7)	139 (64.1)	0.084
Overall	215/760 (28.3)	1180/1736 (68.0)	<0.001

ICU, intensive care unit.

The primary outcome was frequency of discussion of each of the eight components of the infection management framework (table 1).

A convenience sample of pre-implementation and post-implementation interprofessional team bedside round discussions was audited (Monday to Friday) by a non-AMS team member (MSS) using an electronic tool (SimpleSurvey, OutSideSoft Solutions, Quebec, Canada). The ICU team was aware of the auditor but not the audit details.

Data were analysed using descriptive statistics.

RESULTS

During the pre-implementation period (May 2017 to October 2017), 95 audits were completed and 217 were completed during post-implementation (October 2017 to August 2018). After implementation there was a statistically significant increase in discussion of each infection management component, except tailoring of antimicrobial therapy (figure 1, table 1).

Phase 4: *Act*: Over the subsequent 18 months, the intervention was scaled up and implemented into three additional ICUs across the institution. Ongoing feedback is provided to each ICU team weekly during

AMS rounds based on appropriateness of antimicrobial prescribing.¹⁰

DISCUSSION

A structured QI redesign of the daily bedside nursing report in the ICU to include 'infection' led to improved interprofessional discussion of key components of infection management.

This low-resource intervention did not disrupt workflow and increased nursing involvement. Seven of the eight components of infection management discussion increased. Decreased discussion of tailoring of therapy after intervention was unexpected; this warrants future investigation.

Few studies have described nursing engagement in AMS despite their recognised role and to our knowledge this is the first description of such an AMS intervention.^{1 5 6} By leveraging key aspects of behaviour change framework; motivation (need to know information for rounds), opportunity (incorporated into existing workflow), and capability (training and inclusion in discussion increases infection-related knowledge), our novel approach was successful in integrating nurses into bedside discussions of infection management.

Our AMS programme has been active in the ICUs of our institutions for greater than 10 years, therefore our intervention was largely independent of AMS guidance, and focused on the process of communication among the ICU interprofessional team. However, it was not without limitations as it was conducted in a single ICU that was already socialised to AMS. Our audit captured data related to the frequency of sharing of information and statement of planned actions, not quality of discussion. It will be important to explore relationships between quality of discussion and appropriateness of antimicrobial prescribing and patient outcomes. A larger, multicentre study with patient-specific outcomes would help understand the generalisability, feasibility and effectiveness of

Figure 1 Decision-making framework component inclusion before intervention and after intervention.

this approach in improving antimicrobial prescribing and infection management in ICUs.

CONCLUSION

Redesigning bedside nurse reporting to include infection is associated with improved infection management discussion.

Twitter Linda Dresser @ASPLDD and Sarah C J Jorgensen @canuckIDpharm

Contributors LD planned the study, and contributed to the conduct of the study, data analysis and writing of the manuscript. LD is responsible for the overall content as a guarantor. MSS contributed to data collection for the study, initial data analysis and drafting of the study report and manuscript. MM contributed to planning of the study, data analysis and manuscript drafts. LJ contributed to the conduct of the study, data analysis and manuscript writing. SCJJ participated in the data analysis and manuscript revisions. SN participated in study planning, data analysis and manuscript writing. CB reviewed the study plan and provided review of manuscript drafts. AMM approved the study plan and provided review of data analysis and contributed to manuscript revisions.

Funding The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

Competing interests None declared.

Patient and public involvement Patients and/or the public were not involved in the design, or conduct, or reporting, or dissemination plans of this research.

Patient consent for publication Not required.

Ethics approval The study received approval from the research ethics board as a QI initiative.

Provenance and peer review Not commissioned; externally peer reviewed.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which

permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

REFERENCES

- 1 Jeffs L, Law MP, Zahradnik M, *et al*. Engaging nurses in optimizing antimicrobial use in ICUs: a qualitative study. *J Nurs Care Qual* 2018;33:173–9.
- 2 Epstein N. Multidisciplinary in-hospital teams improve patient outcomes: a review. *Surg Neurol Int* 2014;5:295–303.
- 3 Carter EJ, Manning ML, Pogorzelska-Maziarz M. Clinical nurse preparation and partnership in antibiotic stewardship programs: national survey findings are a call to action for nurse leaders. *J Nurs Adm* 2019;49:591–5.
- 4 ANA/CDC. *Redefining the antibiotic stewardship team: recommendations from the American nurses Association/Centers for disease control and prevention Workgroup on the role of registered nurses in hospital antibiotic stewardship practices*, 2017.
- 5 Ha D, Forte MB, Nguyen K, *et al*. Outcomes of bedside Nurse-Driven interdisciplinary antibiotic stewardship and infection prevention rounds. *Open Forum Infect Dis* 2017;4:S487–S87.
- 6 Raybardhan S, Kan T, Chung B, *et al*. Nurse Prompting for Prescriber-Led review of antimicrobial use in the critical care unit. *Am J Crit Care* 2020;29:71–6.
- 7 Morris AM, Bai A, Burry L, *et al*. Long-Term effects of phased implementation of antimicrobial stewardship in academic ICUs: 2007–2015. *Crit Care Med* 2019;47:159–66.
- 8 Langley GL, Moen RD, Nolan KM. *The improvement guide: a practical approach to enhancing organizational performance*. 2nd ed. San Francisco: Jossey-Bass Publishers, 2009.
- 9 Michie S, van Stralen MM, West R. The behaviour change wheel: a new method for characterising and designing behaviour change interventions. *Implement Sci* 2011;6:42.
- 10 Dresser LD, Bell CM, Steinberg M, *et al*. Use of a structured panel process to define antimicrobial prescribing appropriateness in critical care. *J Antimicrob Chemother* 2018;73:246–9.